St. Joseph's Healthcare & Hamilton NUCLEAR MEDICINE REQUISITION So Charlton Avenue East Marian Wing, Level 0 Hamilton, ON LBE 4A6 Hours: 0800 – 1630 Tel: 905-521-6095 Fax: 905-521-6018		Address: Phone: Home: _ Date of Birth (vvvv, Health Card #: _ Physician's Nam	Cell: //;/
Gastrointestinal Biliary Gallbladder Gastric Emptying – Solid Gastric Emptying – Solid Gon Medication Off Medication Gon Medication Off Medication Gon Medication Off Medication Gon Medication Off Medication RBC Liver Scan Liver/Spleen Scan Oncology Sentinel Node – Breast Sentinel Node – Melanoma I lodine 131 Whole Body Scan	Respiratory Lung Scan – Ventilation Perfusion Lung Scan – Quantification Infection Gallium White Blood Cell Scan Bone Scans Whole Body Specific Site: Neurological Perfusion (HMPAO)		Genitourinary Renal Scan Renal with Lasix Renal with Captopril Renal Cortical Plasma G.F.R. Miscellaneous Iodine 131 Therapy (amount) Parathyroid Salivary (Parotid) Thyroid Uptake/Scan Other Exams:

RELEVANT CLINICAL HISTORY:

URGENT REPORT REQUESTED	PHYSICIAN'S SIGNATURE	
Ву:	Print:	
Fax/Phone/Pager:	Sign:	

A COMPLETED REQUISITION, INCLUDING ORDERING PHYSICIAN'S SIGNATURE, IS MANDATORY IN ORDER FOR TEST TO BE PERFORMED.




INFORMATION

Please call 905-521-6095 to change or cancel your appointment. (providing at least 48 hours notice)

Public parking is available. There is also street parking (Pay Station) on Charlton Avenue opposite the Marian and Martha buildings.

> Arrive at least 10 minutes before your scheduled appointment.



Nuclear Medicine – Adult Patient Preparation Instructions **Biliary Scan** Nothing to eat or drink for at least 4 hours prior to test. Test duration -90 minutes. No patient preparation required. Two parts to exam. First portion lasts up to 30 minutes. The **Bone Scan** patient returns 2.5-4 hours later for the second portion of imaging, which lasts for 1 hour. Nothing to eat or drink for at least 12 hours prior to test. Liquid emptying test duration -45**Gastric Emptying Study** minutes. Solid emptying test duration -4.5 hours. May be completed for patient either on or off medication, at physician's discretion (please indicate). If patient is directed not to take medication, then they must be off medication for at least 24 hours. E.g. domperidone/ motilium, metoclopramide/reglan, tegaserod/zelnorm. Lung Scan No patient preparation required. Test duration – 1 hour. Patient to drink 2 glasses of water prior to scheduled test (patient can still void). Check with **Renal Scan** physician for discontinuation of any medications prior to appointment. Test duration -60minutes. **Thyroid Uptake and Scan** Two-day test. In almost all cases, anti-thyroid medications must be discontinued for 5 days, and thyroid hormone replacement medication must be discontinued for 4 weeks. First day, a capsule is given, and uptake is measured 4 hours later. Second day, patient returns for a second uptake and imaging (90 minutes). White Blood Cell Scan Two-day test. First day has two parts, which each take 45 minutes with 4-5 hours between each (Indium) part. Patient should be prepared to be available entire day for test. Next day patient returns for imaging (2 hours). Please notify our office if the patient is on dialysis. White Blood Cell Scan Possible two-day test. First day has three parts, which each take 45 minutes, with 3 hours (HMPAO) between each part. Patient should be prepared to be available entire day for test. Patient may return next day for additional imaging and will be notified of this on the first day. Please notify our office if the patient is on dialysis.