


Lymphoma Chemotherapy – CVPR (oral)

Drug	How drug is given	Day drug is given
Cyclophosphamide	pills	1, 2, 3, 4, 5
Vincristine	IV	1
Prednisone	pills	1, 2, 3, 4, 5
Rituxan	IV	1

You will receive 6 to 8 cycles of CVPR. Each cycle is 22 days. Your chemotherapy schedule may be delayed because of side effects.



Rituxan

Rituxan is a monoclonal antibody, a type of chemotherapy that targets cancer cells. Allergic reactions such as fever, chills and a rash can happen while receiving it through your IV. The first IV treatment is given over 4 to 6 hours to reduce the chance of a reaction. If there is no reaction, it can be given over 90 minutes.

Possible side effects

- Nausea and vomiting, take anti-nausea medications as directed
- Hair loss within 21 days
- Prednisone (steroid) side effects, such as mood swings, irritability and difficulties with sleeping
- Low blood counts which leads to infection, fatigue and bleeding
- Constipation
- Numbness and tingling in fingers and toes

Chemotherapy - What you need to know

There will be a decrease in your blood cells. These include white blood cells, red blood cells, hemoglobin and platelets.

• Call if any side effects seem unexpected or extreme.

Primary Team: Doctor: Nurse: Phone: Phone: Fax: You will need a digital thermometer. You can buy one at most drug stores.

If you have any of these signs of an infection call your primary doctor/nurse right away.



- A temperature of 38°C or 100°F or more. This is a fever.
 Do not take Tylenol, Aspirin, ibuprofen or anti-inflammatory medications without checking with your team first. They may cause bleeding or may hide a fever. Call if you need medication for pain.
- A cough productive of coloured phlegm or sputum.
- Red inflamed skin.
- Diarrhea that does not go away.
- Increased shortness of breath.
- Pain or burning with urination.
- Sore throat.
- Watch for signs of shingles: pain, blisters or a skin rash that may look like chicken pox.

What to watch for:

- Unusual bleeding. Call us right away.
- Constipation. Use your laxatives as needed. Do not use a suppository or enema.
- Numbness and tingling in your fingers and toes.

We encourage you to:

- Attend the chemotherapy class and read the chemotherapy book.
- Eat and drink well (at least 2 liters of non-caffeinated beverages a day).
- Practice good hygiene.
- Practice good mouth care.

Possible severe or long term side effects

- Bone thinning/weakening
- Infertility
- Potential future bone marrow damage

Record your experience with each cycle

Cycle	Date	Notes
1		
2		
3		
4		
5		
6		
7		
8		



Juravinski Cancer Centre 699 Concession Street Hamilton, Ontario L8V 5C2 905-387-9495

PD 8041 – 04/2016 dpc/pted/ChemotherapyLymphomaCVPROralJCC-th.docx dt/April 10, 2017