

Contrast Media (eGFR 30 and under)

Date:	 	
Patient's Name: _		
Referring Doctor:	 	

You have just had an injection of contrast media (x-ray dye) to help the radiologist see your blood vessels and organs on a CT (CAT) Scan.

Instructions for going home

1. Do not take any medications that contain metformin for 48 hours after your CT Scan. See the chart below for the names of these medications.

Trade name	Generic name	
Glucophage, Glumetza, Glucophage XR, Fortamet	Metformin	
Kazano	Alogliptin/Metformin	
Invokamet	Canagliflozin/Metformin	
Xigduo XR	Dapagliflozin/Metformin	
Metaglip	Glipizide/Metformin	
Glucovance	Glyburide/Metformin	
Jentaducto	Linagliptin/Metformin	
Acto Plus Met, Acto Plus Met XR	Plioglitazone/Metformin	
Prandimet	Repaglinide/Metformin	
Avandamet	Rosiglitazore/Metformin	
Kombiglyze	Saxagliptin/Metformin	
Janumet, Janumet XR	Sigtagliptin/Metformin	



2. Contact your referring doctor to have follow-up blood work (eGFR) done to check your kidney function.

This blood work must be done in 2 days after your test (3 days if your test was on a Friday, or 4 days if your test was on a Thursday or long weekend).