

Gastrectomy

You will be having surgery to remove part or all of the stomach. This handout will help you learn about the surgery, how to prepare for surgery and your care after surgery.

Surgery can be done in 2 ways: the laparoscopic or open method.

Laparoscopic method

You are put to sleep with a general anesthesia. The surgeon makes several small incisions (cuts) in your abdomen and one larger incision to remove the stomach. The incisions are closed with dissolvable stitches and tape.

The surgery usually takes about 3 to 5 hours.

Plan to be in the hospital for about 3 to 5 days.

Open method

You are put to sleep with a general anesthesia. The surgeon makes an incision on your abdomen (below the rib cage and above the belly button). The incision is closed with stitches or staples and covered with a dressing.

The surgery usually takes about 4 to 6 hours, but may take longer.


Plan to be in the hospital for about 1 week.

Table of contents

Topic	Page
The stomach	3
Preparing before surgery	4
Day of your surgery	4
Going home	7
Call your surgery if you have	8

Research

During your clinic visit and hospital stay you may be asked to take part in a research study. The research will be explained to you. You can decide whether you want to take part or not. If you decide not to take part, your care will not be affected.



Some or all of your stomach will be removed. You will still be able to eat. You may find that you need to eat small amounts of food 4 to 6 times a day. A dietitian is available to talk with you about eating a healthy diet as you recover from surgery. Preparing before surgery

Keep up with your regular activities, exercise and eat a balanced diet. You want to be as strong as you can before you have the surgery. Arrange for someone to help you around the house, and do errands for you when you come home after surgery.

Follow the pre-op checklist and medication instructions that were given to you during your pre-op visit.

Your belly button needs to be cleaned before your surgery. Wash it with soap and warm water the night before your surgery. It is very important to clean the inside of your belly button very well.

Preparing before surgery

Keep up with your regular activities, exercise and eat a balanced diet. You want to be as strong as you can before you have the surgery.

Arrange for someone to help you around the house, and do errands for you when you come home after surgery.

Follow- the pre-op checklist and medication instructions that were given to you during your pre-op visit.

Your belly button needs to be cleaned before your surgery. Wash it with soap and warm water the night before your surgery. It is very important to clean the inside of your belly button very well.

Day of your surgery

On the morning of your surgery you go to Same Day Surgery (SDS) at the Juravinski Hospital. A nurse will admit you and start your intravenous, or IV. You may be given some medications.

You will then go to the Operating Room (OR), either walking or on a stretcher. Your family may go with you and wait in the waiting area outside of the OR.

After surgery

After your surgery, you will go to the Post-anesthesia Care Unit (PACU). A nurse will be with you as you wake up. Your nurse will check your blood pressure, heart rate and temperature (vital signs) and dressing as needed.

From the PACU you will go to a hospital bed.

Along with your IV, you may have:

- A Patient Controlled Analgesia (PCA) pump – where you push a button to give yourself the pain medication.
- An epidural catheter which is when the pain medicine is given through a thin tube into a small space in your lower back.
- A tube called a foley catheter in your bladder to drain urine.
- Tubes or drains near your incision area to drain extra fluid that can build up after surgery.
- To wear special pressure stockings, (TED stockings) or boots (moon boots). These stockings and boots keep blood moving and help prevent blood clots after surgery.

Pain

You will have pain and discomfort after surgery, especially around your incisions. Pain medicine will be given to you on a regular basis to keep you comfortable and your pain under control. As you heal, your pain should lessen each day. Pain medicine can be given in different ways:

- PCA pump or epidural catheter
- in your intravenous, or IV
- pills - when you are able to drink fluids

Most pain medicines can make you constipated. Stool softeners will be given to help prevent this problem.

You may also have stomach discomfort related to gas pains for a few days after surgery. Walking helps to lessen this discomfort.

Subcutaneous injections

After surgery you will need medication to prevent blood clots. This medication is given with a needle under the skin. It is called a subcutaneous injection.

Dressing and incision

You can shower when you get home. Do not take a bath until your incision is healed.

With the laparoscopic method, each incision should be a dry closed line. Your incisions may be covered with tape (steri-strips), try to keep the tape clean and dry. If the tape falls off, you can leave it off. The incisions will be closed with dissolvable stitches.

With the open method, the incision will be covered by a dressing. The dressing will be removed 2 to 3 days after surgery. The incision will be closed with dissolvable stitches or staples (clips) which will be taken out about a week after your surgery.

If the staples are still in when you leave the hospital, they may be taken out by your family doctor, home care nurse or at your follow-up appointment with your surgeon.

Activity

While in the hospital you will feel tired and unwell. While rest is important, getting up and walking can help most patients feel better and recover quicker. You will be encouraged to do deep breathing, coughing and leg exercises after surgery.

Walking and increasing your activities helps to keep your lungs healthy, prevent blood clots and get your bowels moving.

Move as much as possible while in the hospital:

- Sit up near the edge of the bed and dangle your legs.
 - Sit up in a chair for meals.
 - Sit up when visiting.
 - Walk around the hallway (before doing this, ask for assistance from your nurse or therapist).
-

Feeling tired

It is normal to feel tired after surgery. It may take weeks to months for your energy to return to normal. Everyday do a bit more, walk a little farther.

At home, gradually resume your normal activities. Avoid heavy lifting (5 lbs or 2 kgs), straining or strenuous exercises for at least 6 to 8 weeks. Do not do these until you check with your surgeon.

Eating and drinking

You will have an IV for fluids and medicine. You will have the IV until you are ready to take fluids by mouth. Slowly, you will start drinking clear fluids and progress to solid foods. You may find that you need to eat small amounts of food 4 to 6 times a day. A dietitian is available to talk with you about eating a healthy diet as you recover from surgery.

You may need to have a feeding tube inserted into your stomach to help with your nutrition after surgery.

It may be a few days before you start to pass gas and/or have a bowel movement. You may also feel nauseated or sick to your stomach. Medicine will be given to decrease this feeling.

Going home

You will need transportation home. If you are taking strong pain medicine with a narcotic, such as morphine, do not drive, operate heavy machinery or drink alcohol.

When you go home, you will be given:

- ✓ prescriptions for pain medicine and a stool softener. If not all of your stomach was removed, you will be given a prescription for an antacid.
- ✓ a follow-up appointment with your surgeon.

Your nurse will review with you what you need to know about how to care for yourself at home. Discuss your concerns with your nurse. Ask the nurse to explain or clarify anything you do not understand. We want to ensure you know what to do when you go home.

Call your surgeon if you notice ANY of these problems:

-
- | | |
|--------------|---|
| Fever | <ul style="list-style-type: none">• Your temperature is 38°C (100°F) or higher. |
|--------------|---|
-
- | | |
|-----------------|--|
| Incision | <ul style="list-style-type: none">• Pain in your incision that does not get better with medication.• Your incision is coming open, bleeding, draining or has yellow, green or smelly discharge.• The skin around your incision is red or swollen.• You are concerned about your incision. |
|-----------------|--|
-
- | | |
|----------------|---|
| Abdomen | <ul style="list-style-type: none">• You are constipated, or no bowel movement in 3 days.• You are vomiting or have diarrhea.• You have pain in your abdomen or feel sick to your stomach. |
|----------------|---|
-
- | | |
|--------------|---|
| Other | <ul style="list-style-type: none">❗ You have pain or tenderness in your leg (thigh or calf) with swelling, redness or warmth.❗ You have trouble breathing, chest pain or cough up blood.❗ You have lightheadedness or dizziness that does not go away.❗ You have a rapid heart beat (palpitations), facial flushing, sweating. |
|--------------|---|
-

If you have concerns about your surgery, do not wait for your follow-up appointment, call your surgeon. If you are unable to reach the surgeon, go to Emergency.



**Problems marked with this sign are emergencies.
Call 911 or go to Emergency.**
