

## Care after knee surgery

## For Dr. Punthakee's patients

You had this surgery today:		
☐ knee arthroscopy	☐ lateral release	☐ ACL reconstruction
medial meniscectomy	☐ articular debridement	☐ Fulkerson
lateral meniscectomy	microfracture	hardware removal
dother procedure(s):		

You may remove your dressing/bandage 24 hours after your surgery, and you may then take a shower. Cover your knee with saran wrap. Pat dry after shower. Apply fresh sterile dry gauze and secure with skin tape. Leave the steristrips (the white tape over your incision) on for 10 days after your surgery. Do not take a bath or go swimming for 10 days after your surgery.

## Tonight after your surgery

- take your pain medication as prescribed
- raise your limb
- apply ice to your limb often
- use your crutches or cane for support
- if the bandage is too tight, loosen it
- follow the instructions given by Dr. Punthakee at your pre-op visit

Go to the Emergency Department if you have:

- calf pain
- shortness of breath
- chest pain

These symptoms may represent a blood clot.

You need to be s	seen for a follow-up appointment in about 2 to 3 weeks:	
at Dr. Puntl	thakee's office – 905-577-6677, or	
at the Fract	cture Clinic at the Juravinski Hospital – 905-527-4322, e	xt. 42293.
	ysiotherapy until you have seen Dr. Punthakee, and ha	ave
Your weight bea	aring status and/or use of crutches is as f	ollows:
weight bear	aring as tolerated/crutches for support	
partial weig	ght bearing for weeks	
☐ touch weight	ht bearing for weeks	
□ non-weight	t bearing for weeks	
☐ Remove Zi	immer splint and replace it with your brace in day	/S
Please follow ins	structions given by Dr. Punthakee before your surgery.	
Right	Left	

Please remember to bring this sheet to your appointment at the Fracture Clinic or Dr. Punthakee's office.