

McMaster Children's Hospital Green Book

For Residents and Clinical Clerks 2023-2024



MacPeds

Training the next generation of pediatricians

TABLE of CONTENTS

Welcome to MacPeds!	1
Paging	2
EPIC Tip Sheet - SmartTexts	3
EPIC Tip Sheet - Handover	5
EPIC Tip Sheet – Collect Sexual Orientation and Gender Identity	9
CTU Daily Schedule/Expectations	14
CTU Weekly Schedule	16
CTU Family-Centred Rounds Template	17
MacPeds Handover guide – I-PASS	19
I-PASS	21
GENERAL PEDIATRICS CLINICAL RESOURCES	
Pediatric History & Physical Examination	22
Adolescent Interviews (SSHADESS)	29
Substance Use Disorder, Recognition, and Treatment, M-SSTEP Algorithm	32
Child & Youth Poverty Tool: Hamilton Region	34
Fluid management in children/composition and ordering	36
2 Bag System for DKA	44
Maltreatment Reporting Flow Chart	45
Understanding Your Professional Duty to Report Suspected Child Maltreatment	46
Pronouncing Death	48
Mandatory Infectious Disease Consultations	52
Antibiotic Guide for Common Infections	55
Blood Culture Guidelines – volumes by weight	60
MIS-C Algorithm	62
Reportable Diseases to Public Health	67
Pediatric Proton Pump Inhibitors (PPI) comparison chart	69
ACUTE CARE ALGORITHMS	
HPEWS	70
PACE criteria	71
PALS Algorithms	72

Welcome to MacPeds!

This handbook was designed for the large number of learners and staff from a variety of programs that rotate through McMaster Children's Hospital during their training. It may also be helpful for clinical clerks during their time on the pediatric wards.

Hopefully this demystifies some of the 'pediatric specific' logistics. This is intended only to act as a guideline for general pediatrics use.

We would very much appreciate any feedback, suggestions or contributions emailed to peded@mcmaster.ca.

Sincerely,
Bojana Babic and MacPeds Editors

Permission to copy and distribute this document is granted provided that (1) the copyright and permission notices appear on all reproductions; (2) use of the document is for non-commercial, educational, and scientific purposes only; and (3) the document is not modified in any way. This work is licensed under the Creative Commons Attribution 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Paging

To page someone from within the hospital:

1. dial 87
2. enter the person's pager number (4 digits)
3. enter the call-back extension (5 digits)
4. enter priority code (☐ * then 1 for CODE/STAT, 2 for ROUTINE, 3 for ANYTIME, 4 denotes PHYSICIAN paging)

If you don't know their pager #, wish to leave a typed message, or to wait on an outside line: call **x76443**.

To inactivate/activate your own pager:

1. dial 88
2. enter your own pager #
3. dial 08

Pediatric CTU notes and Templates

SmartTexts

Save these templates as SmartPhrases to edit them, or add them to your favorites as-is.

Progress Notes	
HHS IP PED PROGRESS NOTE	HHS AMB PED COMPLEX CARE PLAN [23313] (to be used by Ped Complex Care in the Complex Care Emergency Action Plan section – not as a progress note)
HHS AMB COMPLEX CARE PLAN NOTE [21762] (to be used by Ped Complex Care for interdisciplinary group note whenever the Complex Care Plan is updated)	HHS AMB PED COMPLEX CARE FOLLOW-UP CLINIC NOTE [22120] (to be used by Ped Complex Care for interdisciplinary group note)
HHS AMB PED COMPLEX CARE INITIAL CLINIC NOTE [22117] (to be used by Ped Complex Care for interdisciplinary group note)	HHS AMB PED COMPLEX CARE VIRTUAL CLINIC NOTE [22121] (to be used by Ped Complex Care for interdisciplinary group note)

Consults	
HHS IP PED CONSULT NOTE	

Discharge	
HHS IP PED DISCHARGE SUMMARY	

Handoff Template	
HHS IP GEN HANDOFF SUMMARY	HHS IP HANDOFF TO DO
HHS HANDOFF SECTION TEXT	HHS HANDOFF PATIENT TEXT (PRINT - LONGER)
HHS HANDOFF PATIENT TEXT (PRINT - BRIEF)	

SmartLinks

Consider adding these SmartLinks within SmartPhrases that you create.

Helpful SmartLinks	
.SCRHRVIS: pull in the data from the hearing and vision screening	.BMI
.SSHADESS: pull in the SHHADESS assessment	

Orders

Order Sets

Save your own versions and edit them, or add them to your favorites as-is.

Order Sets	
GEN PED Admission: Kawasaki Disease or MIS-C	GEN PED Admission: Asthma
GEN PED Admission: Community-Acquired Pneumonia (CAP) (3 months of age and older)	GEN PED Admission: General
GEN PED Admission: Meningitis (3 months and older)	GEN PED Admission: Urinary Tract Infection
GEN PED Admission: Bronchiolitis	GEN PED Discharge to Home or Self Care

Provider Teams

Provider Team System Lists

There are system lists created for your provider teams. Consider saving these lists in a My List for easier access to your teams patients.


General Pediatric Teams	
General Pediatrics Acute Team 1 -MM	General Pediatrics Acute Team 2 - MM
General Pediatrics Complex Care Team - MM	General Pediatrics Consult and Surge Team - MM

Handover


There are multiple tools within Epic to assist in handoff between providers. These tools function both within Patient Lists activity, and in an individual patient charts. This provides functionality for provider teams by combining all individual handover notes into one central list for every member of the provider team.

Writing a handoff report from Patient Lists

1. After selecting a patient, click the **Write Handoff** button pictured below:



2. A template will populate into the sidebar. With this activity, a physician can specify their discipline, note summary details about a patient, and specify 'To Do' items for the receiving provider:




3. Pressing the next button will automatically save the note, and move on to the next patient on the list for documentation on their handover. A provider can alternatively press close to just save the note and close the report.


Adding the 'To Do' column to your Patient List

If you would like to see To Do items from the handoff report directly on your patient list, you can! This configuration can be achieved from any of your My Lists:


1. Right-Click on the My List you would like to add the column to, then select *Properties*.



2. Within the *General* tab, use the Search text field to find the 'To Do' column:



3. Double-click the column from the *Available Columns* list. It will then add the selected column to the *Selected Columns* list and default the column to the right-most position on the patient list. This position is customizable using the arrows below the *Selected Columns* list.
 - a. Note: You can utilize this screen to remove or adjust the positioning of other columns on your My List. The order you set them will match the order they display in the printed list.
4. The column will then appear, representing the To Do items from the handoff report:



Viewing the handoff report from Patient Lists

- To view a handoff report from Patient Lists, single click on the patient to expand the patient snapshot reports.
- Click on the **Handoff** report (pictured below), and see that handoff notes are organized by the provider's specialty. The content of the specialty-specific handoff notes may vary per clinical specialty.
 - If the Handoff report is not available for quick select, expand report options with the magnifying glass!

The screenshot shows the Oncology Patient Lists interface. A list of patients is displayed, including Beacon, Aladdin (26 y.o. M), Beacon, Nadia (7 y.o. F), Beacon, Nicholas (63 y.o. M), Beg, Chandler (20 y.o. M), and Cobble, Jane (32 y.o. F). The 'Beacon, Aladdin' patient is selected, and the 'Handoff' report is highlighted in the left sidebar. The main content area displays the handoff report for Beacon, Aladdin, including patient information, allergies, and a list of providers.

Printing a handoff report

- To print out all handoff reports associated with a patient list, select the **Print** option from the right corner of the top toolbar. Within the dropdown select **Handoff**, to print Handoff reports:

The screenshot shows the Print dropdown menu in the Oncology Patient Lists interface. The 'Print' option is selected, and the 'Handoff' option is highlighted in the dropdown menu.

- An example of the printout is included below:

The screenshot shows the printed handoff report for Beacon, Aladdin. The report is organized into a grid with columns for Patient Information, Allergies, and a list of providers.

Writing a handoff from a patient's chart

1. Within a patient's chart, a provider can select the Handoff activity in the sidebar. A template will populate into the sidebar. This sidebar functions identically to the one found from the Patient Lists activity. A physician can specify their discipline, note summary details about a patient, and specify 'To Do' items for the receiving provider:

The screenshot displays the Epic EMR interface for a patient named Charlie-IM Celaeno. The 'Handoff' sidebar is open on the right, showing a summary of the patient's current status and a list of 'To Do' items. The main chart area shows various medical orders, including medications, scheduled treatments, PRN orders, diet and nutrition, and provider orders. The sidebar also includes a 'Disposition' section for the receiving provider to specify their discipline and note summary details.

Handoff Sidebar:

- Summary:** HPI: Relevant Phob: Patient Active Problem List: Diabetes type 2, controlled. Active medical issues with hypertension.
- To Do:** 1, 2, 3, 4.
- Disposition:** [Empty box for provider notes]

Medical Orders:

- Expiring Medications:**
 - decrease 5% and sodium chloride 0.9% infusion: 100 mL/hr, intravenous, Continuous, Starting on Sun 8/5/22 at 0740. For 2 days. Expires in 20 hours 23 minutes.
 - pantoprazole (Panto IV) 80 mg in sodium chloride 0.9 % 250 mL (0.32 mg/mL) infusion: 8 mg/IV (25 mL/mL), intravenous, Continuous, Starting on Sun 8/5/22 at 0640. For 1 day 13 hours. Expires in 10 hours 23 minutes.
- Scheduled:**
 - erythromycin ophthalmic ointment: Right Eye, Nightly, First dose on Sun 8/5/22 at 2100.
 - furosemide (Lasix) tablet 20 mg: 20 mg, oral, Daily, First dose on Sun 8/5/22 at 0915.
- PRN:**
 - ondansetron (Zofran) 2 mg/mL injection 4 mg: 4 mg, intravenous, Every 8 hours PRN, nausea, vomiting, Starting on Sun 8/5/22 at 1625.
- Diet and Nutrition:**
 - Adult NPO diet: Routine, Diet effective now, Starting on Sun 8/5/22 at 0915, Until Specified.
- Notify Provider Orders:**
 - Notify physician (specify): Routine, Until discontinued, Starting on Sun 8/5/22 at 0840, Until Specified.

Clinical Staff - Collect Sexual Orientation and Gender Identity


Update a patient's name.....	11
Ask about sexuality, gender identity, sex assigned at birth, and pronouns.....	12
Ask about sexuality	12
Ask about gender identity.....	12
Ask about sex assigned at birth	12
Ask about pronouns	12
Respond to patient concerns	13

Collecting sexual orientation and gender identity information from patients allows your organization to monitor trends in care that are impacted by those factors. It also helps to ensure that patients receive appropriate care for risks they might face or organs they have present. Having easy access to information like the gender and name that the patient goes by also facilitates positive interactions between providers and patients.

Use this script to help start conversations with your patients about these topics. Feel free to adapt these scripts as needed, for example, you might have pediatric patients who identify as transgender, but aren't ready to talk about their sexual orientation. We recommend having these conversations with patients somewhere private without parents, guardians, or partners present, because the patient might not feel ready, or safe, disclosing this information in front of others.

Update a patient's name

If a patient goes by a name other than their legal one, record the name so that staff members interacting with the patient see what the patient wants to be called.

1. Ask: "Do you use a different name than your legal name that I should refer to you by?"
2. Click the patient's name from Storyboard.
3. Click  next to the patient's name to open Name Edit.
4. Enter the name by which the patient should be addressed by in the **Preferred name** field and press **Tab**.
 - When you add the patient's preferred name, the **Preferred type** field defaults to First name, Preferred. If you're recording a complete preferred name, update the **Preferred type** field to reflect that.
5. Save your changes.
 - The patient's preferred name now appears on all patient documentation. The patient's legal name still appears on guarantor accounts and coverages, and both names appear on the Interactive Face Sheet.

Ask about sexuality, gender identity, sex assigned at birth, and pronouns

Collect sexual orientation, gender identity, pronouns, and other relevant information by accessing the SOGI SmartForm.

1. Click the patient's gender identity from Storyboard.
2. For guidance on how to collect this information from the patient, use the scripts below.

Ask about sexuality

1. Tell the patient: "To make sure all patients get the best possible care, we would like you to tell us about your sexuality and gender identity. Your answers are confidential, and only visible to those with access to your medical record."
2. Ask: "Which of the following best describes your sexuality? You can choose more than one answer."
3. Read all of the options to the patient.
4. Record the answer or answers in the **Patient's sexual orientation** field. If their answer is not an option, select **Something else**.

Ask about gender identity

1. Ask: "Which of the following best describes your gender identity?"
2. Read all of the options to the patient.
3. Record the answer or answers in the **Patient's gender identity** field. If their answer is not an option, select **Other**.

Ask about sex assigned at birth

1. Ask: "Which of the following best describes the sex that you were assigned at birth? This is the sex that someone observed when you were born, and likely was on your original birth certificate."
2. Read all of the options to the patient.
3. Record the answer in the **Patient's sex assigned at birth** field. If their answer is not an option, select **Uncertain**.

Ask about pronouns

1. Tell the patient: "I use [insert your pronouns here]. What pronouns do you use?"
2. Record the answer in the **Patient pronouns** field.

Respond to patient concerns

You might encounter patients who feel uncomfortable providing their sexuality and gender identity. It's important to be sympathetic to their perspective, because for some patients, disclosing their sexuality or gender might have had negative repercussions in the past.

The table below contains recommended answer to some potential patient responses.

Patient Response	Staff Response	What to Record
"Why do you need to know my sex assigned at birth?"	"It's important for me to have the complete picture so that I can accurately diagnose diseases and suggest the appropriate preventative care for you."	Whatever option the patient ultimately selects, or Choose not to disclose if they decline to answer.
"Can't you tell by looking at me?"	"I'm trained not to make assumptions so that I can record the information accurately. Would you like to hear the options again?"	Whatever option the patient ultimately selects, or Choose not to disclose if they decline to answer.
"It's not your business."	"I understand. I'll record that you don't wish to share."	Choose not to disclose.
"What are pronouns?"	"Pronouns are the words used to refer to a person without using their name. Common pronouns include he/him/his, she/her/hers, and they/them/theirs."	Whatever option the patient ultimately selects, or Decline to answer if they decline to answer.

Division of Pediatric Medicine – CTU 1 and 2 Expectations

Orientation:

At the beginning of each block, the attending should meet with their team members to review the objectives, expectation and schedule of the rotation. The senior resident and/or general pediatric fellow may have valuable input during this time.

Morning Handover:

Morning handover starts at 7:15 or 7:35. Team 1 will be late handover on odd days of the month. Team 2 will be late handover on even days of the month. At 8am the attending will meet with the NP and all trainees to review and divide the patients (10 min). A 20 minute morning report style teaching on one of the admitted patients will occur after. It is therefore important to complete a succinct handover within your team's allotted 20 minutes.

On weekends, morning handover takes place at 8:30 for both teams.

Morning Huddle:

Morning huddle occurs daily to discuss anticipated discharges as well as anticipated length of stay of all patients. These will occur with the attending pediatrician/fellow and senior pediatric resident (SPR) from 9:15-9:30am in 3Z conference room along with nursing and allied health staff. Discharge planning should always be occurring and the team should be aware of potential discharges each day. The attending and SPR should aim to assess and discharge those patients promptly before the start of ward rounds.

See Patients:

After teaching, learners will see their assigned patients. The chart and nursing notes should be reviewed to identify any issues that have arisen over night. The patient should be seen and examined. All lab work and radiological procedures that are pending should be reviewed. The house staff should then come up with a plan for the day and be ready to present that patient during ward rounds. It is not necessary that full notes be written at this time, as there will be time allotted for that later in the day. Potential consults for patients should be discussed with the attending and called during this time to facilitate timely assessments by subspecialty services.

Ward Rounds:

Ward rounds are to take place from 10:00-12:30. During ward rounds the attending paediatrician, SPR, and house staff will round on patients for their team. These are family-centered rounds. Effort should be made to have the family present, either at the bedside or outside the room, while the team is discussing the patient status and management plan. These are also work rounds and orders should be written while rounding on each patient. Some spontaneous teaching during rounds and at the bedside can occur during this time, however there is allotted time for that later in the day.

Multidisciplinary Rounds:

Multidisciplinary rounds take place for senior team members at 1pm on Monday Wednesday and Friday in the 3Z conference room. When there is a long weekend MDR will occur the Tuesday after the long weekend at 1pm in the 3Z conference room. All team members are welcome to attend as interested, but not mandatory.

Patient Care:

During this afternoon residents will follow through with decisions made during ward rounds. This may include arranging investigations, consulting other services, or following up on results. Progress notes, dictations, and other documentation should be completed during this time. Team 1 and 2 members might be called upon to help the Consult admitting in the afternoon if there is a high number of consults.

Afternoon Teaching Sessions:

Afternoon teaching will take place at 2:00pm. Division of General Pediatric Rounds will occur bi monthly on Tuesdays from 12:30-1:30. Please refer to the CTU teaching schedule for locations – this will be posted online as well as on the wards.

Evaluations:

Time is left in the schedule for evaluations. This would be the time to give residents mid-way evaluations, as well as end of rotation evaluations.

Evening Handover:

Evening handover occurs at 16:40 or 17:00. The day team should provide the night team with printed patient lists. The team will then run the list and handover to the on-call team in iPASS format.

Division of Pediatric Medicine, CTU 1, 2, and 5 Weekly Schedule

	Monday	Tuesday	Wednesday	Thursday	Friday
7:15 - 8:30	Resident/Fellow Handover	Resident/Fellow Handover	Resident/Fellow Handover	Resident/Fellow Handover	Resident/Fellow Handover
7:30 - 8:30	Staff Handover	Staff Handover	Staff Handover	Staff Handover	Staff Handover
8:00 - 8:10	*Staff and Team meet to review and distribute patients	*Staff and Team meet to review and distribute patients	*Staff and Team meet to review and distribute patients	*Staff and Team meet to review and distribute patients	*Staff and Team meet to review and distribute patients
8:10 - 8:30	*Staff and Team Morning Report (Teaching) Pediatric Learner Led	*Staff and Team Morning Report (Teaching) Pediatric Learner Led	*Staff and Team Morning Report (Teaching) Staff/Fellow Led	*Staff and Team Morning Report (Teaching) Pediatric Learner Led	*Staff and Team Morning Report (Teaching) Staff/Fellow Led
8:30 - 10:00	Team to see patients, call subspecialists for consults	Team to see patients, call subspecialists for consults	Team to see patients, call subspecialists for consults	Team to see patients, call subspecialists for consults	Team to see patients, call subspecialists for consults
9:15 - 9:30	*Discharge Rounds	*Discharge Rounds	*Discharge Rounds	*Discharge Rounds	*Discharge Rounds
9:00 - 9:15	CQI	CQI	CQI	CQI	CQI
10:00 - 12:30	Team Rounds	Team Rounds	Team Rounds	Team Rounds	Team Rounds
12:30 - 13:30	Lunch	Lunch	Lunch	12:30 - 13:30 Lunch	Lunch
13:00 - 13:30	**Team 1 MDR Rounds	APM Rounds - 2nd Tuesday DPM Rounds - 4th Tuesday	**Team 1 MDR Rounds	*Academic Meeting - 1st Thursday **Clinical Meeting - 3rd Thursday	**Team 1 MDR Rounds
13:30 - 14:00	**Team 2 MDR Rounds	Patient care, follow up with subspecialists regarding consults, Team 1 and 2 support consults	**Team 2 MDR Rounds	Patient care, follow up with subspecialists regarding consults, Team 1 and 2 support consults	**Team 2 MDR Rounds
14:00 - 15:00	*Teaching @20	ASP Rounds	*Teaching @20	*Teaching @20	*Teaching @20
15:00 - 16:20	Patient care, follow up with subspecialists regarding consults, Team 1 and 2 support consults	Patient care, follow up with subspecialists regarding consults, Team 1 and 2 support consults	Patient care, follow up with subspecialists regarding consults, Team 1 and 2 support consults	Patient care, follow up with subspecialists regarding consults, Team 1 and 2 support consults	Patient care, follow up with subspecialists regarding consults, Team 1 and 2 support consults
16:20 - 17:20	Evening Handover	Evening Handover	Evening Handover	Evening Handover	Evening Handover

*TEAM 1 TOUCH BASE AND MORNING REPORT WILL BE IN THE 3C CONFERENCE ROOM
 *TEAM 2 TOUCH BASE AND MORNING REPORT WILL BE IN THE 3Z CONFERENCE ROOM
 *Teaching @20 will be facilitated by Attending and Nurse Practitioners
 *The CTU teaching schedule (along with this schedule) can be found on:
 Medportal -> Podgrad -> Program -> Core Pediatric Residency Program -> Rotations -> Goals/Objectives... -> CTU

MDR: Multi-Disciplinary Rounds
 CTU: Core Teaching Unit Rounds
 DPM: Divisional Pediatric Medicine Rounds
 ASP: Rounds: Antibiotic Stewardship Programme Rounds

Version 4.0.0.2

Rounding Process: 3C – McMaster Children's Hospital

Purpose of rounds is to:

- collaboratively develop and communicate a plan of action for each patient with the Interprofessional health care team, the patient and the patient's family
- facilitate safe and timely patient discharge planning
- provide a forum for education
- provide excellent patient care

Pre-Rounding Agenda

(to be done before round start time of 10:00)

Team Member	Pre-Round Tasks
SPR/Fellow/Staff/NP	<ul style="list-style-type: none"> • See watchers and new admissions as needed • See patients for pre-round discharge • Call urgent consultants and arrange urgent investigations/procedures
Resident/Clerk/NP	<ul style="list-style-type: none"> • Talk to patient/family and bedside nurse • Examine patient • Review physician progress notes, interprofessional team notes, medication profile, flow sheet data (VS, I/O, weight) and new investigation results • Obtain computer for rounds
Bedside RN	<ul style="list-style-type: none"> • Gather pertinent data to be presented (weight change, I/O, Urine Output, Fluid Balance, VS, etc)
Charge RN	<ul style="list-style-type: none"> • Create list showing patient rooms for each nursing assignment • Help with coverage while bedside nurse attends rounds

Notes for Rounds:

- Rounding time goal: **(10:00-12:30)**
- **Team 1 starts on 3C, Team 2 Starts on 3Z** unless watcher needs to be seen first
- SPR/Staff to monitor time and lead brief teaching points (1 per patient)
- Once a bedside RN joins rounds aim to round on all of their assigned patients
- Rounds to be completed in patient room unless otherwise requested by patient/family
- Information to be presented in a sensitive manner for all patients/families
- Pharmacist will attend rounds with each team on alternate days to assist in reviewing of medications

Post-Rounding Agenda

Team Member	Afternoon Tasks
SPR/Fellow/Staff/NP	<ul style="list-style-type: none"> • Help facilitate and arrange consults, investigations, or procedures • Complete family update if not already done at rounds (include junior learner if possible) • Ensure learners attend afternoon teaching on time • Ensure Handoff list is updated and review iPASS with learners before evening handover
Resident/Clerk/NP	<ul style="list-style-type: none"> • Arrange investigations/consults as discussed at rounds • Follow-up on any outstanding investigations • Attend afternoon teaching • Complete progress notes, update problem list and orders if there are any changes post rounds • If done early, help with new admissions/consults or transfers • Ensure Handoff list is updated with clear plan for oncoming night-time team • Communicate any changes to the plan made at rounds with bedside nurse/family <i>Note: Patient care to be completed before evening handover</i>
Bedside RN	<ul style="list-style-type: none"> • Implement orders
Charge RN	<ul style="list-style-type: none"> • Ongoing updates on plan from bedside RN and NP as applicable
Business Clerk	<ul style="list-style-type: none"> • Fax forms/consults for LHIN and Community providers and upload into Media folder on EPIC • Print Stickers for referral forms as needed

Rounding Process: 3C – McMaster Children's Hospital
Rounding Template

Section	Details	Presenter
ID	Brief sentence: gender, age, reason for admission and pertinent PMHx Note: Only discuss presentation and initial workup if pt is new to team Does this patient have a POST?	Resident/ Clerk/NP
Prioritized Issue List:	1. Issue #1 2. Issue #2 ...	Resident/ Clerk/NP
Note: Nurse to present information below but resident/clerk/NP to review prior to rounds.		
Systems Review <i>Focus on concerns or pertinent negatives</i>	VS & HPEWS Values for most recent VS & HPEWS score & team to visually review VS trends on WOW (**no longer saying VSS) <i>Reassess frequency of VS & continuous monitoring daily</i>	Bedside Nurse
	CNS: LOC, Pain Management, Withdrawal (WAT Scores), NVS	
	CVS: Perfusion, IV access (difficult?), IVF & Rate, Consider TKVO?	
	RESP: Cough, Breath Sounds, WOB details, FiO ₂ , Secretions, Suctioning, NS Drops, Incentive Spirometry, Chest Tube	
	GI/GU: Diet, Feed Amount (Route), TPN, Last BM (Consistency), G/JT?, Ostomy?, Drains? Abdo Girth, Emesis, Today's Wt (↓ or ↑ from previous) , Foley or I/O Cath? Urine Output (ml/kg/h), Fluid Balance	
	MSK/Skin: Ambulation/Mobility, Pressure Sore Risk, Breakdown, Rashes, Wounds/Dressings	
	ID: Isolation, New Diarrhea/Vomiting/Cough	
	Social: Patient/Family Questions & Concerns, Stressors/Coping, Safety Concerns, BSR	Pharmacist will join rounds when available
	Investigation /Procedures: Frequency of Bloodwork, Outstanding Investigations or Procedures	
	Medications Review MAR, PRNs Given, Held/Refused , Stop Dates	
Which Disciplines are Currently Involved → <input type="checkbox"/> CLS <input type="checkbox"/> OT <input type="checkbox"/> PT <input type="checkbox"/> RD <input type="checkbox"/> RT <input type="checkbox"/> SLP <input type="checkbox"/> SW <input type="checkbox"/> Wound Nurse <input type="checkbox"/> Other		
Assessment and Plan:	<ul style="list-style-type: none"> Summary of relevant information from Systems Review, Physical Exam and Investigations/procedures. Review subspecialists involved Present active issue list with assessment & plan for each <ol style="list-style-type: none"> Issue #1 – Assessment (i.e. cause of issue/differential) & Plan (further investigations, reassess meds, consult services, etc) Issue #2 – Assessment & Plan ... Consider → Hydration, Growth and Nutrition, and Discharge Planning → If Home & Community Care Support Services – HCCSS (formerly LHIN) services are needed: Make sure to order in EPIC & give forms/order to Business clerk to E-fax. Reassess Daily → CRM/Cont SPO₂, Accurate I/O, freq of VS/Wt/bloodwork Orders → review all active orders, learner/NP who is not presenting on current patient to update orders before moving on to next patient Review and Update Problem List 	Resident/ Clerk/NP
Conclusion	Final Summary (+/- revisions) of Plan	SPR/Fellow NP/Staff
Outstanding Questions and Concerns		Pt, Family & Team
Post Rounding Multi-D Rounds (MDR)	Mon-Wed-Fri → Team 1 at 13:00 & Team 2 at 13:20 Meet in the 3Z Conference Room with the Multidisciplinary Team to run through the patient lists.	NP/Staff

AM Handover Guidelines

7:15 am Early Team Handover

The Early Team will receive handover at this time

- ☐ Team 1: Even Days
- ☐ Team 2: Odd Days



Please bring your own printed list to handover

**Weekend & Holiday
AM Handover
starts at 8:30am**

The overnight JRs (junior residents) & clinical clerks will present new patients

Spent 2 to 3 min for each patient and discuss:

- ☐ Name, age, main presenting complaint(s)
- ☐ Brief history with the most important pertinent positives/negatives
- ☐ Relevant past medical history
- ☐ Brief summary of objective findings (physical exam, investigations)
- ☐ Admitting diagnosis and plan



Try to remember to focus on information that will change or inform patient management!

**Subspecialty
AM Handover
occurs in the
PICU at either
7:30am or
9:00am on
weekdays and
at 9:30am on
weekends &
holidays**

JRs present team issues:

- ☐ Briefly state overnight issue(s) and management
- ☐ Inform the team of any issues that need follow-up or task(s) that were handed over the night before
- ☐ If there are no overnight issues or follow-up, simply state "No issues" or skip the patient and move on



7:35 am Teaching Session

Clinical Clerk/JR/SR will present a case seen overnight or a topic of interest. Points to include:

- ☐ Salient clinical features
- ☐ Diagnosis and differential diagnosis for the patient
- ☐ Acute treatment options and brief long-term management goals (evidence-based, if possible)



7:35 am Late Team Handover

The Late Team will receive handover at this time:

- "Team on Take = Handover Late"**
- ☐ Team 1: Late Handover on Odd Days
- ☐ Team 2: Late Handover on Even days



7:50 am Heme-Onc & Team 3 Handover

Heme-Onc & Team 3 will handover at this time to incoming residents, fellows or staff

PM Handover Guidelines

*CTU Seniors are expected to contact the Weekend Day SPR to handover the weekend plans for patients on their respective teams

4:30 pm



Team 3

The incoming team will print their own lists – please have them updated by 4:30pm



- ☐ Team 3 will give handover to the covering JPR (junior pediatric resident) along with the SPR (senior pediatric resident)
- ☐ Note: If this handover is expected to take longer than 10 minutes, the JPR will accept the rest of handover outside of the room and Team 1 or 2 will start handover



4:40 pm
Early Team Handover

The outgoing team will present team handover
Please follow the IPASS format



I Illness Severity	Stable, "watcher," unstable
P Patient Summary	<ul style="list-style-type: none"> • Summary statement • Events leading up to admission • Hospital course • Ongoing assessment • Plan
A Action List	<ul style="list-style-type: none"> • To do list • Time line and ownership
S Situation Awareness and Contingency Planning	<ul style="list-style-type: none"> • Know what's going on • Plan for what might happen
S Synthesis by Receiver	<ul style="list-style-type: none"> • Receiver summarizes what was heard • Asks questions • Restates key action/to do items

(Starmer et. al, 2012)

Subspecialty
PM Handover
occurs at **5:30pm**
in the PICU on
weekdays, weekends
and holidays



5:00 pm
Late Team Handover

The **Late Team** will give handover at this time:

"Team on Take = Handover Late"

- ☐ Team 1: Late Handover on **Odd** Days
- ☐ Team 2: Late Handover on **Even** days

Note: If the early team arrives late for handover, or has exceeded the allotted handover time, their handover will be interrupted by the Late Team Handover at 5:00pm. The Early Team can then resume handover once the Late Team has finished

5:20 pm
Heme-Onc Handover

Heme-Onc will handover to the JPR & SPR at this time.
Please ensure that patient lists are updated.

I	Illness Severity	<ul style="list-style-type: none"> • Stable, “watcher,” unstable
P	Patient Summary	<ul style="list-style-type: none"> • Summary statement • Events leading up to admission • Hospital course • Ongoing assessment • Plan
A	Action List	<ul style="list-style-type: none"> • To do list • Time line and ownership
S	Situation Awareness and Contingency Planning	<ul style="list-style-type: none"> • Know what’s going on • Plan for what might happen
S	Synthesis by Receiver	<ul style="list-style-type: none"> • Receiver summarizes what was heard • Asks questions • Restates key action/to do items

FIGURE 1
 Elements of the I-PASS mnemonic.

Handover Format – the I-PASS breakdown

- I:** Status: stable vs. watcher
- P:** One-line summary of child and reason for admission
 List of active issues +/- relevant management
- A:** Overnight action list
- S:** Anticipated overnight issues with management plans
- S:** Brief clarification from receiver (1-2 questions) if needed. If further questions, defer to end

PEDIATRIC HISTORY & PHYSICAL EXAMINATION

HISTORY

Identifying Data:

- Name, sex, age (years + months), race, who accompanies child, significant PMHx

Chief Complaint: in patient's or parent's words

History of Presenting Illness (HPI):

- Open-ended questions, allow parents/child to express their concerns
- Similar HPI details to an adult history
- Establish time line: "when was your child last well?", "what happened next?" etc.
- Select key symptoms and expand:
 - colour, character, quantity of vomit etc,
 - OPQRST of pain, aggravating/relieving factors etc
- Always ask about recent exposures to ill contacts – family, school

Past Medical History (PMHx):

- Significant ongoing medical problems
- Prenatal history:
 - Mother's age, gravida, live births, abortions etc
 - Planned vs unplanned pregnancy, onset of prenatal care
 - Complications, smoking, drinking, meds, drug use in pregnancy
 - Gestational age at birth
- Birth history:
 - Spontaneous vs induced labour, duration, complications
 - Presentation: breech, vertex, transverse
 - Interventions required: forceps, vacuum, c-section
 - Resuscitation required, Apgars, birth weight (conversion chart)
 - NICU, Level 2 nursery admission, duration
- Newborn history:
 - Common problems: jaundice, poor feeding, difficulty breathing
- Hospitalizations and significant accidents
- Surgical history

Medications – including dose changes, compliance

Allergies – list specific reaction

*** Immunizations** – ask specifically about Prevnar, Menjugate, Varivax, Synagis (if neonate).

Feeding History (if relevant):

- Breast feeding: exclusively?, duration, frequency
- Formula: brand, how is it prepared/diluted, # of feedings/day, quantity
- Solids: when started, tolerated, any reactions
- Vitamins (especially iron and Vit D): which ones, how often, dose
- Present diet: cereals, fruit, vegs, eggs, meat, amt of cow's milk
- Any difficulties with feeding? Any concerns from primary physician about poor weight gain?

Developmental Milestones (if relevant):

- “Have you ever had any concerns about your child's development?”
- “How does child compare with siblings/peers?”
- Ask about current milestones in each category as appropriate for their age:
 - Gross motor
 - Fine motor, vision
 - Speech, hearing
 - Social skills
- Use major milestones (walking, first word, toilet training, etc) to assess previous development (*Reference in Development Section*)
- Use Denver II charts etc. to assess current stage of development

Social History

- Who lives at home? Who are primary caregivers? Parents work outside the home?
- Does the child attend daycare? How many other children? In a home vs. institution?
- Stability of support network: relationship stability, frequent moves, major events (death in family etc), financial problems, substance abuse in the home
- Has CAS ever been involved?
- School adjustment, behaviour problems, habits (nail-biting, thumb sucking etc), sleep changes

- How has this disease affected your child/ your family?
- What does your family do for fun? What does your child do for fun?
- For an asthma history: smoke, pets, carpets, allergens in the home, family history of asthma / atopy.

Family History:

- Are parents both alive and well? How many siblings? Are they healthy?
- Are there any childhood diseases in the family?
- Consanguinity – are mother and father related in any way?
- Relevant family history (3 generations) – autoimmune hx in Type I DM, atopic hx in asthma etc
- Draw pedigree if possible for genetic assessment

Review of Systems:

General: feeding, sleeping, growing, energy level

Signs of illness in kids: *activity, appetite, attitude* (3 A's)

HEENT: infections (how often, fever, duration): otitis, nasal discharge, colds, sore throats, coughs, nosebleeds, swollen glands, coughing or choking with feeding

Cardio:

Infants: fatigue/sweating during feedings, cyanosis, apneas/bradycardic episodes

Older kids: syncope, murmurs, palpitations, exercise intolerance

Resp: cough, wheezing, croup, snoring, respiratory infections

GI: appetite, weight gain (growth chart), nausea/vomiting, bowel habits, abdominal pains

GU: urinary: pain/frequency/urgency, sexually active, menarche/menses, discharge/pruritis/STDs

MSK: weakness, sensory changes, myalgias, arthralgias, 'growing pains'

Neuro: headaches, seizures (febrile vs afebrile, onset, frequency, type), tics, staring spells, head trauma

Skin: rashes, petechiae, jaundice, infection, birthmarks

PHYSICAL EXAMINATION

General Inspection

- Sick vs not sick?
- Toxic appearance? listlessness, agitation, failure to recognize parents, inadequate circulation (cool extremities; weak, rapid pulse; poor capillary refill; cyanotic, gray, or mottled colour), respiratory distress, purpura
- Level of consciousness
- Nutritional status – well nourished?
- Developmental status (“pulling up to stand in crib”, “running around room”)
- Dysmorphic features – look specifically at face, ears, hands, feet, genitalia

Vital Signs:

- Include Temperature, Heart Rate, Respiratory Rate, Blood Pressure and O₂ saturation

NORMAL PEDIATRIC VITAL SIGNS

Age	HR	SBP	RR
Newborn (<1 wk)	120-160	60-70	30-60
Neonate (<1 mos)	120-160	75-90	30-60
Infant (<1 year)	110-140	75-120	20-40
Preschool (3-5yrs)	90-120	75-125	20-25
Child (6-12 yrs)	80-110	83-120	16-24
Adolescent (>12 y)	70-100	90-130	12-18
Adult (>18 yrs)	60-100	90-130	12-18

Anthropometrics (plot on growth curves at every visit!):

- Height (supine length to 2 years, then standing height)
- Weight
- Head circumference (generally birth to 2 years, >2 yrs if specific concerns)
- Plot BMI (kg/m^2) on updated CDC growth curves for appropriate BMI for age

Hydration Status

- Comment on mucous membranes, tears, skin turgor, sunken eyes, in addition to appropriateness of vital signs, etc.
- For classification of mild, moderate, severe dehydration – see “Fluids & Electrolytes”

HEENT:

- Head: dysmorphic features, shape of skull, head circumference, fontanels in infants
- Eyes: strabismus, pupillary response, funduscopy, red reflex in infants, conjunctivitis
- Ears & pharynx exam in any child with a fever!
- Nose: turbinates, deviation of septum, presence of polyps?
- Mouth: lips (lesions, colour), mucous membranes including gingiva, tongue, hard/soft palate,
- Dentition: presence of teeth, tooth decay
- Neck: lymphadenopathy, palpation of thyroid, webbing (Noonan, Turner syndrome), torticollis

Cardiovascular:

- HR, BP, apical beat, heaves/thrills
- Perfusion:
 - o Pulses – strength/quality, femoral pulses in all infants
 - o Capillary refill time
 - o Skin colour: pink, central/peripheral cyanosis, mottling, pallor
- S1/S2, extra heart sounds (S3, S4)
- Murmurs:
 - o Timing (systole, diastole, continuous)
 - o Location of maximal intensity, radiation
 - o Pitch and quality (machinery, vibratory, etc),
 - o Loudness (I – VI / VI)

Respiratory:

- Audible stridor, sturor, wheeze, snoring
- Position of child, ability to handle secretions
- Signs of distress: nasal flaring, tracheal tug, indrawing
- RR, O₂ saturation (current FiO₂), level of distress
- Able to speak in full sentences (if age appropriate)
- Depth and rhythm of respiration
- Chest wall deformities: kyphosis, scoliosis, pectus excavatum/carinatum
- Finger clubbing

Abdomen:

- For peritoneal signs: ask child to jump up and down or wiggle hips, to distend and retract abdomen “blow up your belly and then suck it in”
- Inspection: scaphoid/distended, umbilical hernias, diastasis recti
- Auscultation: presence of bowel sounds
- Percussion: ascites, liver span, Traube's space for splenomegaly
- Palpation: hepatosplenomegaly?, tenderness, guarding (voluntary, involuntary), masses (particularly stool presence in LLQ)
- Stigmata of liver disease: jaundice, pruritis, bruising/bleeding, palmar erythema, caput medusa, telangiectasia, ascites, hepatosplenomegaly

Genito-urinary:

- Anal position, external inspection (digital rectal examination in kids ONLY with clinical indication), Sexual Maturity Rating
- Male infants: both testes descended, hypospadias, inguinal hernias
- Females: labia majora/minora, vaginal discharge, erythema/excoriation of vulvovaginitis (NO speculum exam if pre-pubertal), Hymenal exam if indicated.

MSK:

- Gait assessment, flat feet vs toe walking vs normal foot arches
- Standing: genu valgum “knock knee” vs genu varum “bow legged”
- Joints: erythema, swelling, position, active/passive range of motion, strength, muscle symmetry
- Back: kyphosis, scoliosis

Neurological:

- Overall developmental assessment
 - o Try playing ball with younger children, or even peek-a-boo!
- Level of consciousness (Glasgow Coma Scale if appropriate)
- Newborns: primitive reflexes, moving all limbs, presence of fisting?
- Cranial nerves: by observation in infants, formal testing in older children
- Motor: strength, tone, deep tendon reflexes, coordination
- Sensory: touch, temperature, position/vibration sense
- Cerebellar: gait (heel to toe, on heels, on toes, finger-to-nose, rapid alternating movements in older children, Romberg (eyes open then closed)

Derm:

- Jaundice, pallor, mottling, petechiae/purpura
- Rashes, birthmarks, hemangiomas, stigmata of neurocutaneous disorders

For helpful physical exam videos: <http://learnpediatrics.com/videos/>

ADOLESCENT INTERVIEWING (SSHADESS Screen)

- Interview teens alone with parents invited to join at the end (Alternatively, you can start with the parents in the room and have them leave at some point)
- Allow adequate, uninterrupted time to inquire about all aspects of their life, and high-risk behaviours in private setting
- Assure **confidentiality** at beginning of interview, and prior to discussing drug use and sexuality. Remember caveats of confidentiality (ie. if you are at risk of harm to yourself or others, or if someone is hurting you)
- Remember to obtain routine history including: Past Medical History, Meds, Allergies and Vaccines (HPV, hepatitis, meningococcal in particular)

Strengths

- What do you like doing?
- How would you describe yourself? How would your best friends describe you?
- Tell me what you are most proud of.

School

- Name of school, grade level
- What do you enjoy most/least about school?
- How many days have you missed or arrived late to school?
- How are your grades? Any different from last year?
- Do you feel like you are doing your best at school? (If no) Why not?
- What would you like to do when you get older?

Home

- Tell me what home is like...
- Who lives at home? How does everyone get along? What do you argue about? What are the rules like at home?

- Family members – ages, occupations/education, health status, substance abuse

ADOLESCENT INTERVIEWING (Continued)

Activities

- What do you do for fun? On weekends?
- Do you feel you have enough friends? Who are your best friends? What do you do together?
- Do you have any extra-curricular activities?

Drugs/Substance Use

- Have you ever tried cigarettes? Alcohol? Marijuana?
- Do you drink alcohol? Binge drinking on weekends?
- For younger teens: ask about friends' use and peer pressure
- Cover all drug classes: hallucinogens, amphetamines, rave drugs, IV drugs, crack cocaine, OTC meds, anabolic steroids. If D positive see M-SSTEP for next steps to screen for risk of withdrawal.
- What age did you start? Frequency of use? How much?
- What do you like/dislike about X? Why do you use X ?
- Do you use alone? Any police involvement? Dealing?

Emotions/Eating/Depression

- Have you been feeling stressed? Do you feel nervous a lot?
- Do people get on your nerves more than they used to?
- Have you been having trouble sleeping lately?
- Do you have concerns about your weight/shape?
- Have you tried to change your weight/shape in any way?
- Any bingeing or purging behaviours (includes diuretics/laxatives)
- Tell me what you eat/drink in an average day...
- *****TIP:** Use growth curves to estimate 'healthy weight' based on height*
- Have you been feeling down, sad, or depressed?
- Depression screen – SIGECAPS

- Have you thought of hurting yourself or someone else? Have you ever tried to hurt yourself?

ADOLESCENT INTERVIEWING (Continued)

Sexuality

- Are you attracted to anyone? Tell me about that person.
(Using gender-neutral language)
- Are you attracted to guys, girls, or both?
- What kind of things have you done sexually? Kissing?
Touching? Oral Sex? Have you ever had sexual intercourse?
- How many sexual partners have you had?
- What do you use for contraception/STI prevention (condoms, OCP, Depo-provera, Emergency Contraception etc.)
- Any history of sexually transmitted infections?
- Have you ever been pregnant or gotten someone pregnant?
- Have you ever been forced or pressured into having sex?

Safety

- Do you regularly use: seatbelts? Bike helmets? Appropriate gear when snowboarding/skateboarding or other sports?
- Do you feel safe at school? Have you ever been bullied?
- Does anyone at home own a gun?
- Have you ever been the victim of violence at home, in your neighbourhood or at school?
- Has anyone ever hurt you or touched you in a way that was hurtful or inappropriate

M-SSTEP Algorithm



Critical Reminders

- ▶ Risk of withdrawal warrants **EARLY** assessment and planning at admission.
- ▶ Withdrawal can be life threatening and can begin within hours of last use.
- ▶ Abrupt stops in substance use or periods of abstinence can initiate withdrawal. Patients with prior history of withdrawal are more likely to experience withdrawal. **Know your STEP-A answers!**
- ▶ Unintentional/unknown polysubstance use (Fentanyl lacing) is common. Youth are often unaware of this risk and may not disclose polysubstance history.
- ▶ Abrupt stops/period of abstinence lowers tolerance and this increases risks of poisoning and death. At discharge, or AMA consider role of Naloxone education (see M-SSTEP Resource Guide for Naloxone information).
- ▶ In benzodiazepine withdrawal, avoid using antipsychotic medication due to the risk of lowered seizure threshold
- ▶ In opioid withdrawal, avoid benzodiazepines due to the added risk of respiratory depression; and avoid Clonidine due to false lowering of the patient's opioid requirements without preserving tolerance.

Positive Drug Use on SHADES?

NO

YES
Assess Withdrawal Risk

Provide positive reinforcement and education on substance use

STEP A - Assess Acute Withdrawal Risk

1. Assess risk of acute withdrawal from alcohol, opioids and benzodiazepines, by asking detailed substance history. Use "Step-A" questions.
 - S** - What **substances** are being used and how (i.e. smoking, injecting, oral, snorting)
 - T** - What was the **timing** of last use?
 - E** - Have they had **experience** with overdose or withdrawal?
 - P** - What is the **pattern** of use (i.e. daily, weekly, monthly, binge)?
 - A** - What **amount** is being used (i.e. quantity)?
2. Assess for current withdrawal signs and symptoms (from alcohol, opioids and benzodiazepines) using appropriate withdrawal assessments tools:
 - Alcohol (CIWA-Ar)
 - Opioids (COWS)
 - Benzodiazepines (CIWA-B)
3. Check urine (order UDRUGCOMP) if STEP A questions indicate use of opioids and/or benzodiazepines. Consider adding serum ethanol, ASA, Acetaminophen. Inform patient and obtain consent.
See M-SSTEP Clinical Resource Guide for additional support

Current symptoms of withdrawal?
= Crisis Presentation

No current symptoms of withdrawal but possibility of future withdrawal?
= Incidental Withdrawal

No current symptoms of withdrawal and no perceived risk for future incidental withdrawal?

Withdrawal plan needed NOW

STEP B - Assess Acute Withdrawal Risk

1. With data from the STEP A questions and the MSSTEP resource guide, develop a withdrawal plan. Ensure MRP is aware.
2. Communicate withdrawal risk (crisis or incidental) to front-line RN.
3. Consult Addiction Medicine Team (AMT) (via HHS paging, available Mon-Sun 0800-1700), for help with pharmacological management of acute withdrawal.
4. Consult pediatric social worker to provide supportive counselling as part of the admission order set.
5. Consider psychiatry consult if needed for assistance with the co-management of psychiatric issues
6. Consider nicotine replacement therapy (if applicable). See HHS policy and HHS order set

1. Continue to monitor vitals. If abnormal HPEWS or development of withdrawal symptoms, reconsider if patient is in crisis presentation and need for withdrawal plan

2. Use M-SSTEP Clinical Resource Guide for help
3. Notify SPR & MRP for further assessment

STEP C and Step D
(see page 2)

STEP C – Optimization of Withdrawal Plan

1. If not already done, consult AMT. Consider consulting again if optimization is needed.
2. Consider nicotine replacement therapy (if applicable) to ease withdrawal symptoms.
 - a. See HHS policy and HHS order set
3. Consult Child Life Specialist to provide support re hospitalization and coping strategies.
4. Consult pediatric social worker (if not already done) for supportive counselling and support during hospitalization.
5. Consult Adolescent Medicine (during daytime hours) for:
 - a. Any youth presenting with withdrawal (either crisis or incidental).
 - b. Any youth with substance use and co-existing chronic illness (i.e. diabetes, cystic fibrosis)
6. Consult Psychiatry Consult Liaison (Mon-Fri 0800-1600) to assist with withdrawal plan, mental health issues.
7. Screen patient for the presence of a substance use disorder (within 24 hours of admission). Use the CRAFFT Screening Tool for Adolescent Substance Abuse



STEP D – Discharge Treatment Planning

1. Ensure appropriate follow up referrals are made *
 - Rapid Access Addictions Medicine Clinic (RAAM): for ongoing withdrawal management and/or diagnostic assessment for presence of a substance use disorder
 - Alternatives for Youth (AY): substance use and/or disorder support; family resources
 - Young Adult Substance Use Program (YA-SUP): Youth 17+ only for substance use assessment and treatment; concurrent disorder treatment; family support
2. Provide education about Naloxone Kits and where to access kits
3. Provide resources – pamphlets, resource list*

*Refer to M-SSTEP Resource for more information (Found here)

M-SSTEP (McMaster's Substance Support for Teens Through Education and Partnership)

Child & Youth Poverty Tool: Hamilton Region

A practical tool for clinicians



By Orianna Mak, MD & Ania Van Meer, MD
Updated February 2022

Poverty poses a significant risk to child and youth health, and should be addressed as such.

What can we do as healthcare providers to address this risk factor and reduce inequities?

ASK...

1

Do you have trouble making ends meet?

2

Do you have trouble feeding your family?

3

Do you receive the child tax benefit?

4

Do you have legal or immigration challenges?

5

Do you have a safe and clean place to live?

6

Are you/your child in need of dental care?

7

Are you concerned about your/your child's mental health?

TURN OVER FOR
RESOURCES



1. Do you have trouble making ends meet?

Benefits & Supports

<https://benefitswayfinder.org/>

Special Support Program & Affordable Transit Pass

<https://www.hamilton.ca/social-services/support-programs>

Income Tax Clinics

<https://hamiltontaxhelp.ca>

2. Do you have trouble feeding your family?

Food Banks

Search "food banks" at <https://redbook.hpl.ca/>

Good Food Box

Affordable fresh fruit & veg program - on hold due to COVID

https://www.environmenthamilton.org/good_food_box

City of Hamilton Food Access Guide

Information on free/low-cost meals, student nutrition programs, food cooperatives, & more

<http://foodaccessguide.ca/>

Student Nutrition Programs

View their "Impact Report" for a list of program locations

<http://www.tastebudshamilton.ca>

Children's Breakfast Club

Nutritious, hot breakfast for children & families every school day from 7:45-8:45am at Compass Community Health (438 Hughson St N).

Community Fridges

24/7 access to free food in a community fridge & freezer

- 44 Greendale Dr, Hamilton, ON, L9C 5Z4
- 249 John St N, Hamilton, ON, L8L 4P4
- 204 Ottawa St N, Hamilton, ON, L8H 3Z5
- Corner of Locke St & Stanley Ave

Essential Aid - Infant & Toddler Food Bank

100 Main St E, Suite 201, Hamilton, ON, L8N 3W4

Mon/Wed/Fri 10am-12pm, Tues/Thurs 7pm-9pm

3. Do you receive child & family benefits?

Tax-payment to eligible families to help with the cost of raising children <18. Includes the **Canada Child Benefit**, **Ontario Child Benefit**, & **Child Disability Benefit**

<https://www.canada.ca/en/revenue-agency/services/child-family-benefits.html>

Child Care Subsidy

Financial support for child care through the City of Hamilton

<https://www.hamilton.ca/social-services>

4. Do you have legal or immigration challenges?

Hamilton Community Legal Clinic

(905) 527-4572, <https://hamiltonjustice.ca/>

Ontario Council of Agencies Serving Immigrants

Information & services for newcomers <https://settlement.org/>

Community Legal Education Ontario <https://www.cleo.on.ca/en>

Legal Aid Ontario

Legal aid services for those who financially qualify. Call 1-800-668-8258 (Mon-Fri 8am-7:30pm) or visit <http://www.legalaid.on.ca/>

Hamilton Immigration Partnership Council

<https://www.hamiltonimmigration.ca/>

Justice For Children & Youth

Legal services for youth under 18 and homeless youth under 25

<https://jfcy.org/en/>

5. Do you have a safe and clean place to live?

Hamilton Housing Help Centre

(905) 526-8100 www.housinghelphelpcentre.ca

Ontario Renovates Program

Financial assistance for low-income families to repair substandard housing to a minimum level of health & safety

<https://www.hamilton.ca/social-services/support-programs/ontario-renovates-program-homeowners>

Good Shepherd Family Centre

24-hour intake line (905) 528-9442

Notre Dame House & Community Outreach Services

Youth-only (16-21) shelter & services (905) 308-8090

6. In need of dental care?

Healthy Smiles Ontario

Free dental care for children 17 & younger from low-income households. Automatic enrolment for families on OW/ODSP.

<https://ontario.ca/healthsmiles>

Dental Health Bus

Mobile outreach clinic offering free emergency dental services

<https://www.hamilton.ca/public-health/clinics-services/dental-health-bus>

Public Health Services Dental

Currently closed <https://www.hamilton.ca/public-health/clinics-services/public-health-services-childrens-preventive-dental-clinic>

For urgent dental needs, call (905) 546-2424 ext. 5369

Further list of dental programs & services in Hamilton

<https://chs.hwcdsb.ca/support/parentresources/?fileID=172209>

7. Mental health concerns?

Crisis Outreach & Support Team (COAST) 24/7 (905) 972-8338

Trans Lifeline 24/7 (877) 330-6366

Contact Hamilton

Entry point for child & youth mental health & developmental services.

Self-ref at (905) 570-8888 or info@contacthamilton.ca

<https://contacthamilton.ca/>

Alternatives for Youth

Free substance use & mental health services (905) 527-4469

<https://ay.on.ca/>

Canadian Mental Health Association - Hamilton

(905) 521-0090, <https://cmhahamilton.ca/>

Free mental health phone apps for youth

Clear Fear, Smiling Minds

FLUID MANAGEMENT IN CHILDREN

Children are at high risk of dehydration:

- Higher % total body water compared to adults
- Higher body surface area : mass ratio
- Higher metabolic rates
- Higher insensible losses
- Limited access to free water

Management of Dehydration

1. Assess severity and type of dehydration
2. Deficit Replacement
3. Maintenance Fluids
4. Replace Ongoing Losses
5. Reassessment and Monitoring

1. Assess Severity and Type of Dehydration

- Severity of dehydration dictates urgency of situation and need for acute resuscitation
- Degree of dehydration represents the percentage of body weight lost due to acute loss of fluids and electrolytes
- Degree of dehydration estimated based on history and physical exam (See Table on next page)
- Type of dehydration reflects relative net losses of water and electrolytes – based on serum Na⁺ or osmolality

Type of Dehydration	Electrolyte Status	Clinical Features
Hypotonic or Hyponatremic	Serum Na ⁺ < 130 mEq/L Serum Osm < 270	Exacerbated signs of dehydration Risk of seizure
Isotonic or Isonatremic	Serum Na ⁺ 130-150 mEq/L Serum Osm 270 – 300	
Hypertonic or Hypernatremic	Serum Na ⁺ > 150 mEq/L Serum Osm > 300	Decreased signs of dehydration Irritable, increased tone and reflexes

Assessment of Degree of Dehydration

	Mild	Moderate	Severe
% Weight Loss (by age)	5% (< 1 year) 3% (> 1 year)	10% (< 1 year) 6% (> 1 year)	15% (< 1 year) 9% (> 1 year)
General Appearance	Alert Thirsty	Drowsy Restless	Lethargic Cold, mottled limbs
Tachycardia	Absent	Present	Present
BP	Normal	Orthostatic Hypotension	Hypotension
Respirations	Normal	Deep +/- rapid	Deep + rapid
Fontanel or Eyes	Normal	Slightly depressed	Sunken
Tears	Present	+/-	Absent
Mucous membranes	Moist	Dry	Very dry
Skin turgor	Normal	Reduced	Tenting
Cap Refill	Normal	>2 secs	>>2 secs
Pulses	Present	Weak	Not palpable
Urine output	Normal	Oliguria	Anuria

2. Deficit Replacement

- To calculate fluid deficit:

$$\text{Fluid Deficit} = \% \text{ Dehydration} \times 10 \times \text{body weight}$$

- Each 1% dehydration = 10 ml/kg fluid deficit

Oral Rehydration Therapy

- ORT is the first-line treatment for mild - moderate dehydration
- Requires close monitoring and compliance of patient and parents
- Goal is to replace the deficit over 4 – 6 hours and replace ongoing losses by oral intake
- Initial rates of ORT:
 - Mild – 1 mL/kg/5 mins
 - Moderate – 2 mL/kg/5 mins
- Prefer solutions with balanced amounts of sodium and glucose (see table below)
- Feeding should be continued throughout oral rehydration to help maintain gut nutrition

Solution	Glucose (mEq/L)	Na (mEq/L)	K (mEq/L)	Base (mEq/L)	Osmolality
WHO	111	90	20	30	310
Rehydrate	140	75	20	30	310
<i>Pedialyte</i>	140	45	20	30	250
Pediatric Electrolyte	140	45	20	30	250
Infantlyte	70	50	25	30	200
Naturlyte	140	45	21	48	265

Parenteral Therapy (IV)

- IV therapy indicated for severe dehydration and patients who fail ORT due to: vomiting, refusal, or difficulty keeping up with losses
- Preferable site is IV, if unable to start IV use IO

(i) Restore Intravascular Volume

- Goal: expand ECF volume to prevent or treat shock and maintain perfusion

IV Bolus 10 – 20 ml/kg of N/S or RL
run over 15-20 mins or rapid push

- NEVER use hypotonic solution for boluses
- Avoid dextrose-containing solutions
- Monitor for improvement following each bolus – assess HR, BP, mental status, etc.
- May repeat boluses until patient is hemodynamically stable – If unstable, call Peds 1000!

(ii) Ongoing Deficit Replacement

- Goal: replace remainder of fluid deficit over next 24 hours
- Subtract boluses from deficit calculation
- Replace $\frac{1}{2}$ deficit in first 8 hours, second $\frac{1}{2}$ deficit over next 16 hours
- Solution:
 - D5 NS + 20 mEq/L KCL in isotonic dehydration
 - D5 $\frac{1}{2}$ NS + 20 mEq/L KCL in hypernatremic dehydration
- Solution chosen based on type of dehydration and serum electrolytes
- IV fluid rate should include deficit replacement + maintenance fluids (see next section)

3. Maintenance Fluids

- Fluid and electrolyte requirements are directly related to metabolic rate
- All patients, regardless of degree of dehydration, should be considered for maintenance fluids if oral intake is impaired
- Holliday-Segar Rule – maintenance fluid requirements calculated based on body weight for resting hospitalized patients (based on 100 ml for each 100 kcal expended)

Body Wt (kg)	Daily Rate (100-50-20)	Hourly Rate (4-2-1 rule)
The first 10 kg (1-10 kg)	100 mL/kg/day	4 mL/kg/hr
The 2nd 10 kg (11-20 kg)	+ 50 mL/kg/day	+ 2 mL/kg/hr
Any Additional kg (>20 kg)	+ 20 mL/kg/day	+ 1 mL/kg/hr

- Insensible water losses = cutaneous + pulmonary water losses which are calculated as $\sim 300 - 500 \text{ cc/m}^2$
- Important to assess factors affecting insensible and/or urinary fluid losses – may need higher maintenance rate
- Normal Na^+ and K^+ requirements 2 – 4 mEq/kg/day
- Also affect factors affecting Na and K balance – may need to include additional supplementation
- Solution:
 - D5 $\frac{1}{2}$ NS + 20 mEq/L KCL
 - D5 NS + 20 mEq/L KCL
- Adding 5% dextrose to maintenance solution prevents protein catabolism (Use D10W in neonates and hypoglycemia)
- Solution chosen based on type of dehydration and serum electrolytes
- D5 $\frac{1}{2}$ NS + 20 mEq/L KCl provides 4 mEq/100 mL Na^+ and 2 mEq/100 mL K^+
- Only add K^+ if patient is voiding

4. Replace Ongoing Losses

- Assess patient for additional fluid losses – diarrhea, vomiting, polyuria, drains, etc
- Estimate output over 4-6 hours then replace volume
- Replacement fluid dependent on source of losses

Replace...	With...
Gastric Losses (Vomiting)	$\frac{1}{2}$ NS + 10 – 20 mEq/L KCl
Stool or Intestinal losses (Diarrhea)	Add HCO_3^- to $\frac{1}{2}$ NS + 10 – 20 mEq/L KCl
CSF losses	0.9% NS
Urine Output	As indicated
Losses due to Burns	Increase fluid administration (Parkland formula)

5. Reassessment and Monitoring

- Important to continually assess patient's hydration status and fluid requirements
- Monitor HR, BP, Cap refill, mental status and urine output
- Accurate INS and OUTS, repeat weight measurements
- May require cardiorespiratory monitor, CVP, ECG
- Check serum electrolytes routinely while patient on maintenance fluids
- Other labs as indicated: BUN, Cr, serum osmolality, urine specific gravity, urine osmolality
- Adjust type and rate of IV fluids depending on clinical and biochemical indicators of volume status
- Discontinue IV fluids once patient has returned to normal status and tolerating normal feeding

Comparison of IV Solutions

IV Solution	Na ⁺ (mEq/L)	K ⁺ (mEq/L)	Cl ⁻ (mEq/L)	Dextrose (g/L)	Osmolarity (mOsm/L)
Sodium Chloride 0.45%	77			0	154
Sodium Chloride 0.9% (0.9 NaCl, NS)	154		154	0	308
Sodium Chloride 3%	513			0	1030
Dextrose 5%	0			50	250
Dextrose 5% Sodium Chloride 0.2%* (D5 0.2NS)	39			50	320
Dextrose 5% Sodium Chloride 0.45% (D5 ½NS)	77		77	50	405
Dextrose 5 % Sodium Chloride 0.9%	154			50	560
Dextrose 10%	0			100	505
Dextrose 10% Sodium Chloride 0.2%*	39			100	575
Dextrose 10% Sodium Chloride 0.45%*	77			100	660
Dextrose 10% Sodium Chloride 0.9%*	154			100	813
Dextrose 3.3% Sodium Chloride 0.3% (½ * ⅓)	51		51	33.3	273
Lactated Ringers†	130	4	109	0	273

†Also contains Calcium (Ca^{2+}) 1.5 mmol/L, and Lactate (HCO_3^-) 28 mmol/L

*These solutions are not commercially available

Commonly used solutions are highlighted

Guidelines for Prescribing Maintenance IV Fluids in Children

- These are general guidelines for ordering maintenance IV fluids (IVF) only, and do not apply to resuscitation or complicated fluid and electrolyte disorders. **Seek additional advice/appropriate consultation in the event of fluid and electrolyte abnormalities.**
- Consider IV fluids as DRUGS - individualize prescriptions *daily* according to objectives, and monitor for potential side effects.
- Be aware that the commonest side effect of IVF therapy is HYPONATREMIA, particularly in patients at risk, and if hypotonic solutions are used

Step 1:

Determine IV fluid rate, according to "maintenance fluid" requirements, and replacement of deficit or ongoing losses (Total Fluid Intake (TFI)). In general maintenance fluid rate is calculated by the "4:2:1" guideline, but should be individualized according to the clinical condition and patient assessment

Step 2: The choice of fluid is dependent the individual patient.

Consider **ISOTONIC** IVF for the following patients:

- CNS disorder, Diabetic ketoacidosis
 - Patients at risk of hyponatraemia: acute infection, post-operative patients and burns, Plasma Na < 138
- Add K⁺ to provide 1-2 mEq/kg/day, if patient has urine output
- Add Dextrose to prevent hypoglycaemia/ketosis (exceptions: hyperglycaemia, brain injury)

Consider **HYPOTONIC** IVF for the following patients:

- Patients with an EFW deficit - e.g. hyponatraemia, ongoing EFW losses (renal, GI, skin)
- Patients with established 3rd space overload - e.g CHF, nephrotic syndrome, oliguric renal failure, liver failure
- Limited renal solute handling indicated - e.g. neonatal population, hypertension

*Based on a sodium plus potassium concentration in the aqueous phase of plasma of 154mEq/L, assuming that plasma is 95% water with a plasma sodium of 140 mEq/L and a potassium concentration of 4 mEq/L.

Weight (kg)	ml/hour
0-10	4kg/hour
11-20	40 + (2kghr)
>20	60 + (1kghr)

IV solution	Na (mEq/L)	K (mEq/L)	Cl (mEq/L)	% Electrolyte Free Water (EFW)*
H y p o t o n i c	0.2% NaCl in DSW	34	0	34
	0.45% NaCl in DSW	77	0	77
	Lactated Ringers	130	4	109
	0.9% NaCl in DSW (ISOTONIC)	154	0	154
				0

Step 3: MONITORING while on IV fluid	Measure and record as accurately as possible	Labs:
<p>Clinical status: hydration status, urine output, ongoing losses, pain, vomiting, peripheral edema, and general well-being.</p> <p>Daily weights</p> <p>Reassess TFI indications for and fluid prescription at least every 12 hours.</p>	<p>Fluid balance: must be assessed at least every 12 hours</p> <p>Intake: All IV and oral intake (including medication). Ensure this matches desired TFI.</p> <p>Output: all losses (urine, vomiting, diarrhea etc.)</p>	<p>Serum Electrolytes - at least daily if primary source of intake remains IV, or more frequently depending on clinical course, or in the presence of documented electrolyte abnormality.</p> <p>Urine osmolality/sodium and plasma osmolality as indicated, for determining etiology of hyponatraemia.</p>

2 bag system for DKA

Glucose (mmol/L)	BAG A (No dextrose)		BAG B (dextrose)		Combined Dextrose Concentration
	% of IVF rate	mL/h (fill in below)	% of IVF rate	mL/h (fill in below)	
Greater than 18.0	100%		0%	zero	0%
15-18.0	75%		25%		2.5%
12-14.9	50%		50%		5%
9-11.9	25%		75%		7.5%
Less than 9.0	0%	zero	100%		10%

+

Equals
IVF rate
(D)
mL/h

What to do as a Resident/Fellow if you learn of possible child maltreatment while working at McMaster Children's Hospital or the Ron Joyce Children's Centre

Legend

PG trainee = Resident or Fellow
HCP = Health Care Professional
MRP = Most Responsible Physician
CPA = Child Protection Agency (CAS, FACS etc)
CAAP = Child Advocacy and Assessment Program (the McMaster Child Maltreatment team)

PG trainee has new, **first-hand** information that a child may be at risk of, or has experienced, maltreatment

ASAP, PG trainee notifies most senior PG learner (if applicable) and MRP

Situation requires report to CPA

PG learner and MRP discuss situation with care team and make plan for informing the family (and patient, if appropriate)

Reasons for concern and plan to report to CPA are discussed with the family unless there is a risk that violence may result

PG learner and MRP call CPA for region where the child resides

Date & time of report and discussion with family documented in the chart

Unclear if situation requires report to CPA

Consider requesting guidance from on-call CAAP MD (available 24/7 through paging)

Report required or still unclear if required¹

PG trainee is given information by another HCP that a child may be at risk of, or has experienced, maltreatment

ASAP, PG trainee notifies most senior PG trainee (if applicable) and MRP

MRP and most senior PG trainee assist the HCP with managing their duty to report.

Report not required²

1. You can discuss a case with the CPA without providing the child or family's name. If a report is required you will have to provide this information.
2. If, after consulting, you still feel that you should call CPA, you must do so. No one can lawfully prevent you from calling.

Understanding Your Professional Duty to Report Suspected Child Maltreatment

Neglect: Failing to ensure that one or more of a child's needs are met (e.g. nutrition, safe home environment, supervision, clothing, education) **resulting in harm or a risk of harm**. Medical neglect occurs if the child has a medical, mental or developmental condition that requires treatment and the caregiver does not ensure that the child receives the necessary care. **Neglect makes no reference to what the caregiver's feelings or intentions are.**

Physical Abuse: Any deliberate physical force inflicted on a child by a caregiver that results in pain, injury or creates a genuine risk of harm to the child.

Sexual Abuse: Involvement of a child in any form of sexual activity by a person with a duty of care to the child (e.g. family member, family friend, care provider).

Emotional Abuse: Repeatedly treating a child in a way that negatively impacts their sense of self-worth and self-esteem, such as repeatedly yelling, ignoring, rejecting, demeaning, isolating.

Exposure to Intimate Partner/Family Violence/Conflict: Direct and indirect exposure to violence, threats, verbal abuse etc. between people (usually adults) in the home.

Caregiver Fabricated Illness in a Child: The caregiver actively engages in behaviour which causes the child to undergo unnecessary medical care; such as by providing false information and/or directly causing symptoms in the child.

What are your reporting responsibilities as a Health Care Professional?

The law makes it clear.

Child, Youth and Family Services Act of Ontario

The duty to report applies to children 15 and under. Reporting on behalf of 16 and 17 year-olds is lawful and encouraged, but not legally required.

“Despite the provisions of any other Act” – Your duty to report supersedes all other legislation, including PHIPA

“A person who performs professional...duties with respect to children has reasonable grounds to suspect” that a child “has suffered” or “there is a risk that the child is likely to suffer” any form of maltreatment. – You don't have to be sure. You just have to have a reason to suspect that maltreatment has occurred or may occur.

“the person shall immediately report the suspicion and the information on which it is based to a society.”- You have to manage with the situation when it arises.

“A person who has a duty to report...shall make the report directly to the society and shall not rely on any other person to report on the person's behalf” – The person who acquires the concerning information must be the person to report it to the Child Protection Agency. You must not get someone else to do it for you and nobody can lawfully compel you to do so on their behalf.

However, if a group of clinicians concludes that a report is required (e.g. a team concludes that medical neglect is occurring) one person can be delegated to make the report. This should be the person most capable of explaining the risk of harm.

“A person who has additional reasonable grounds...shall make a further report”
- Every new concern must be reported, even if it happens the same day.

And,

It is unlawful for someone to use their authority to prevent you from making a report.

Please refer to the accompanying flow-chart for guidance on what to do if you become aware of possible child maltreatment while working at McMaster Children's Hospital or RJCC

Pronouncing Death

Before the Death:

Understand the patient's symptom management plan:

- Which medications/dosing to give for which symptoms

Goals of Care:

- What are the family's hopes? (ex: minimizing symptoms, promoting wakefulness)

Communication:

- Any tips about communication strategies to help support the child and family

Understand who is involved and who wants to be notified:

- This will likely include the MRP, QoLA Care and the Senior Resident/Fellow

Before Pronouncing:

Don't rush: The pronouncement does not need to occur at the exact moment death is suspected. Take time to collect your thoughts before entering the room.

Know your patient and setting: Know the name of the child and the family members. Ensure there is adequate seating. The child may be in their parent's arms.

You will need: 1) Death certificate, 2) Death notification checklist, 3) Consent for autopsy, 4) autopsy request (if applicable). These documents can be found in the pediatric ward document cabinet, PICU or NICU.

Pronouncing Death:

Introductions and expectations:

- Take time to briefly introduce yourself if this is your first time meeting the family. Explain your role and that you will be listening to the child's chest for what will seem like a long time. Plan to listen for a period of **two minutes**.

Death Pronunciation Exam:

- Listening for breathing and heartbeat is all that is needed. It is **not** necessary to check pupil reactivity, pain response or peripheral reflexes. If there is any visible respiratory effort, or an audible heart beat, then the child cannot be pronounced dead. You may then stop listening, explain there is still a heart rate and that you will listen again after some time. You can explain that it is common for a child's heart to take time to completely stop.

Language recommendations:

- Make eye contact and use a phrase like, *"Mr/Mrs (Last Name), I am sad to say that (Child's Name) has died."*
- Use the word *"died/dead."* Do not use euphemisms.

Give space:

- Tell the family you will give them some time and that you will check in with them to answer questions and discuss next steps. Some families like to have someone with them after their child has died, provide what support feels right to you.
- Ask the family if there is anyone they would like you to contact on their behalf.

Give anticipatory guidance:

- Explain the next steps and offer to answer any questions. Explain that the family can stay with the child as long as they wish. The child's body can be picked up from the hospital room by the funeral home or be picked up from the morgue. If an autopsy is planned, the child will go to the morgue.
- Ask the family about their wishes for an autopsy.
 - *"We always ask about the option of having an autopsy. This is something we offer to all families whose children have died. Not all families choose this for their child, but some do. Would you like to hear more about what this means?"*
 - *If yes: "An autopsy is usually performed in the event that the cause of death is unknown or if families want more answers about their child's death. Their organs will be examined closely by a physician specially trained to do so. The autopsy leaves a scar on the chest, which can be covered up. Some families worry that their child's autopsy will delay burial or cremation. In our experience, the delay is usually 24 to 48 hours."*

Note: If the autopsy includes the brain, it may take longer.

After Patient Death:

Notify relevant parties:

- Must include: MRP, Next of Kin (if not present), Coroner (if meets requirements), Trillium Gift of Life
- Might include: QoLA Care (if involved), Child Protective Services (if open file), Social Work (if involved), CAAP (if involved), Chaplaincy (if requested)

Submit documentation:

- Place death certificate, death notification checklist +/- autopsy consent and request into manila folder in chart marked *"Death Documents"*. These will be sent to health records.
- Complete a death dictation (see below).
- It's **strongly** encouraged to debrief with the MRP or another member of the care team. This can be after the event, or can be planned for a later date.

Death Dictation:

Use "Discharge Summary" dictation code. Copies to MRP and Family Physician.

"E.g. Admission date: (Date). Date and time of death (Date, Time). Admission diagnosis (Diagnosis). Cause of death: (use cause from death certificate). Other diagnoses at time of death: (List). Course in Hospital: (brief highlight of events surrounding admission and death). The family has been notified of the patient's death. The death packet has been completed. Autopsy was offered to the family and the family has elected to have/not have an autopsy."

Special Considerations:

Coroner's Case: Must call 1-855-299-4100 if one of the following: Cause of death is uncertain, CAS has been involved in the last year, child is a ward of the state, suspected abuse or neglect

Trillium Gift of Life Network (TGL): 416-363-4001

- Call if there is currently an open file or if the family is willing to receive a call from TGL.
- Inform the family that TGL can provide more detailed information on tissue donation.
"E.g. Many families are interested in learning more about tissue donation. If you would like more information, Trillium Gift of Life will connect with you. Trillium Gift of Life is the government organization responsible for coordinating tissue donation services across the province."

Appendix A: Sample Medical Certificate of Death

Cause of Death		
11. Part I		I
Immediate cause of death	(a) Neuroblastoma	Approximate interval between onset and death 3 years
	due to, or as a consequence of	
Antecedent causes, if any,	(b)	
	due to, or as a consequence of	
	(c)	
	due to, or as a consequence of	
Underlying cause of death (Stated last)	(d)	
Part II		II
Other significant conditions contributing to the death but not resulting in the underlying cause given in Part I		

Cause of Death		
11. Part I		I
Immediate cause of death	(a) Aspiration Pneumonia	Approximate interval between onset and death 1 Day
	due to, or as a consequence of	
Antecedent causes, if any,	(b) Cerebral Palsy	5 Years
	due to, or as a consequence of	
	(c)	
	due to, or as a consequence of	
Underlying cause of death (Stated last)	(d)	
Part II		II
Other significant conditions contributing to the death but not resulting in the underlying cause given in Part I	Other diagnoses, not related to death: ex. Diabetes	

Mandatory Infectious Diseases Consultations

As a move towards achieving accreditation goals, and quality indicators the following list of conditions has been developed to help to improve clinical outcomes for rare but severe infections.

ID consultation:

Please call ID for consultation within 24 hours for all patients with the following conditions.

Condition-based

- 1) Any proven meningitis or encephalitis
- 2) Any proven orbital cellulitis or mastoiditis with intracranial extension
- 3) Any suspected/proven bone or joint infection
- 4) Any suspected/proven necrotizing skin infection
- 5) Any suspected/proven endocarditis
- 6) Any severe pneumonia complicated by parapneumonic effusion requiring drainage
- 7) Fluid refractory septic shock requiring admission to PICU with >24hours of persisting end organ dysfunction
- 8) Severe COVID requiring medical intervention beyond dexamethasone

Organism-based:

- 1) Severe *C. difficile* infection (including toxic mega colon, admission to ICU, or significant lab abnormalities)
- 2) *Staphylococcus aureus* bacteremia
- 3) Invasive Candida infection (Candidemia, Candida meningitis, Hepatosplenic candidiasis)
- 4) Any suspected infection with multi-drug resistant pathogens or requiring a carbapenem, such as a patient with a known current or past history of infection or colonization with: ESBL producers, multi-drug resistant *Pseudomonas*, septic patient worsening despite >24 hours of broad spectrum antimicrobials
- 5) Any suspected/proven infection requiring broad spectrum antimicrobials (carbapenems, caspofungin, amphotericin B, voriconazole, posaconazole)
- 6) Complex pathogens requiring specific microbiologic information
- 7) Any suspected/proven malaria
- 8) Any suspected/proven TB (tuberculosis infection)

Microbiology Testing Information

Please refer to EPIC Procedure Catalog

- Stool multiplex PCR (Salmonella, Shigella, Campylobacter, Yersinia and Shiga toxin producing E.coli (STEC) including E.coli H7:O157)
- Respiratory virus PCR (Influenza A & B, RSV, rhinovirus, parainfluenza, human metapneumovirus, adenovirus and COVID-19)
 - a COVID only PCR is also available if recent exposure etc.
 - *Mycoplasma pneumoniae* / *Chlamydia pneumoniae* is an add-on test
- CSF virus PCR (HSV, VZV, enterovirus and parechovirus [< 5 years of age])
- Stool Virology includes PCRs for Adenovirus, Rotavirus, and Norovirus 1 and 2. If you are interested in Enterovirus results this must be ordered separately
- HSV Blood PCR (search altona in EPIC)

Microbiology Tests which require ID or Microbiologist approval:

- Bacterial CSF PCR (Pediatric: *S. pneumoniae*, *N. meningitidis*, *H. flu*, *Listeria*; Neonate: also includes *GBS*, *E. coli* K1)
- Pleural fluid/Empyema PCR: *GAS*, *MSSA/MRSA*, *S. pneumoniae*
- Bone biopsy/joint fluid: *GAS*, *MSSA/MRSA*, *S. pneumoniae*, *Kingella kingae*
- CMV or EBV blood PCR
- 16s bacterial PCR on sterile sites
- 18s fungal PCR
 - *Bacterial and fungal PCR will only be run if culture is negative*

DIAGNOSIS OF URINARY TRACT INFECTIONS IN PEDIATRICS

An appropriately collected urine sample is important for the accurate diagnosis of a urinary tract infection in children. An inadequate sample may lead to overtreatment of what is a contaminated sample, potentially overlooking the real cause of infection in a febrile infant, or failure to diagnose and treat a true urinary tract infection. **Urinalysis – Routine and Culture** is the correct EPIC Order. Culture will not be done unless the Urinalysis is abnormal.

Following a review of national and international guidelines, the following recommendations are to be followed for submitting a urine sample from children for bacteriological culture:

1. DO NOT collect urine in a urine bag, the so-called “bagged urine”. These samples are associated with significant contamination of $>50\%$. This sample source is no longer available to order and will be rejected for culture by the laboratory. Where bacterial contamination is not of concern (e.g. urine for CMV, metabolic screens), a bag urine may be appropriate. Urine collected into a “clean” cotton swab in a “clean” diaper and squeezed out is NEVER an appropriate sample to send for culture.
2. In children who are toilet trained, a “clean catch” urine can be collected. Where possible, start collecting the urine after the first few drops which will wash away any contaminants. Identify the specimen type as “Urine, Clean Catch”.
3. In young infants < 6 months of age, there can be value in attempting to collect a clean catch urine sample by suprapubic cutaneous stimulation, the so-called “Bladder Stimulation/Tap”, or a variation of this method, in a well hydrated infant. If a clean catch urine can be collected within 5-10 minutes

of trying, this sample can be submitted, ensuring that it is identified as “clean catch”. If this is unsuccessful, an “in/out catheter” sample should be collected.

4. In a child who is not toilet-trained or where collecting a timely clean catch urine is difficult, the best sample to collect is using an “in and out catheter” as this minimizes any contamination. The specimen type MUST be correctly identified as an “in and out catheter” so that the appropriate work-up can be done in the laboratory.
5. There will be occasions where there may be other sources of urine for culture, e.g. indwelling catheter, nephrostomy tube. Please ensure the correct specimen type is identified on the order.

Urinalysis is sensitive and specific for the diagnosis of urinary tract infections in children, EVEN IN YOUNG INFANTS. For most infants and children, it is recommended that urine culture is only performed when the urinalysis is positive (leukocyte esterase or nitrites). Culture, however, should be performed on children who are neutropenic, regardless of the urinalysis, and pregnant patients.

In settings where microscopy is clinically appropriate, Urine R and M can still be ordered.

Canadian Pediatric Society. Urinary tract infection in infants and children: Diagnosis and management. Paediatr Child Health 2014;19:315-19

American Academy of Pediatrics. Urinary tract infection: Clinical Practice guidelines for diagnosis and management of the initial UTI in febrile infants and children 2 to 24 months. Pediatrics 2011;128:595-610

Labrosse M, Levy A, Autmizguine J, et al. Evaluation of a New Strategy for Clean-Catch Urine in Infants. Pediatrics. 2016;138(3):e20160573

Tzimenatos L, Mahajan P, Dayan PS et al. Accuracy of the urinalysis for urinary tract infections in febrile infants 60 days and younger. Pediatrics 2018 141(2):e20173068

McDaniel CE, Ralston S, Lucas B, Schroeder AR. Association of diagnostic criteria with urinary tract infection prevalence in Bronchiolitis. JAMA Pediatrics 2018.5091

ANTIBIOTIC GUIDE FOR COMMON INFECTIONS


Infection	Major Organisms	Antibiotic	Duration	Notes
Otitis Media	<i>S. pneumoniae</i> , <i>H. influenzae</i> (non-typable), <i>M. catarrhalis</i> (2-20%) Group A <i>Streptococcus</i> (5%)	Preferred: Amoxicillin PO (75-90mg/kg/DAY divided BID) if type 1 allergy → Clarithromycin PO if non-type 1 → Cefprozil PO OR Ceftriaxone IM OD x 3 days If initial therapy fails: Amoxicillin-clavulanate (Clavulin) PO If type 1 allergy → call ID Amoxicillin-clavulanate IV if no concern about CNS infection	10 days (age < 2) 5 days (age > 2)	watchful waiting appropriate when: - > 6mo - healthy child (NO immunodeficiency or chronic disease) - absence of anatomical abnormality of head and neck - NO Dots, no otalgia, NO history of complicated otitis media - illness not severe - reliable parents CPS statement 2016
Mastoiditis	<i>S. pneumoniae</i> , <i>H. influenzae</i> (non-typable), <i>M. catarrhalis</i> , Group A <i>Streptococcus</i> , <i>Staphylococcus aureus</i> , less commonly <i>anaerobes</i> (<i>fusobacterium</i>)	Ceftriaxone + metronidazole if concern about intracranial spread may need CNS dosing depending on extent of infection If suspect MRSA (e.g. previous colonization) or if severe disease, add vancomycin and involve ID	IV treatment once improving and source control completed (usually ~ 7 days), and then completion of 4 weeks total with PO Amox-clavulanate	MUST INCLUDE CLINICAL findings of mastoiditis = postauricular tenderness, erythema, swelling (loss of crease), fluctuance/mass /discharge, protrusion of auricle (radiologic evidence of mastoid effusion alone IS NOT DIAGNOSTIC) - assess for complications of subperiosteal abscess, abscess of the mastoid, subcutaneous abscess, abscess of the facial nerve, thrombosis - ENT consultation
Orbital cellulitis	Group A <i>Streptococcus</i> , <i>Streptococcus pneumoniae</i> , <i>Staphylococcus aureus</i> , <i>H. influenzae</i> , <i>anaerobes</i>	Oral stepdown: Amoxicillin-clavulanate Amoxicillin-clavulanate IV if no concern about CNS infection Ceftriaxone + metronidazole if concern about intracranial spread may need CNS dosing depending on extent of infection If suspect MRSA (e.g. previous colonization) or if severe disease, add vancomycin and involve ID	Mild orbital cellulitis – usually 2-3 weeks total duration, but will depend on whether there are abscesses and/or bone or CNS involvement.	Mandatory ID consult
Community-acquired pneumonia	3 mo - 4 yrs Viral >> Bacterial (<i>S. pneumoniae</i> , group A <i>Streptococcus</i>) >> Atypicals (<i>Mycoplasma</i> , <i>Chlamydia</i> , <i>Legionella</i>) 5 - 18 yrs <i>S. pneumoniae</i> , atypicals, GAS	Outpatient/admitted to ward: High dose Amoxicillin PO (75-90mg/kg/DAY divided TID (max 1g po TID)) or Ampicillin IV Atypical pneumonia (often seen in generally well older children): Clarithromycin PO Plavural effusion/empyema Ampicillin IV if not getting drained Amoxicillin/clavulanate if chest tube being inserted (pending culture and PCR) Consider Vancomycin if history of MRSA infection in patient or family Admitted to ward/ICU: Acetaminophen Ceftriaxone IM/IV + Vancomycin IV	Mild Nonspecific pneumonia (no admission required): 5 days Pneumonia requiring admission to hospital: 7-10 days Empyema/effusion: consult ID (likely weeks)	Features of atypical pneumonia: subacute onset, non-lobar infiltrate, minimal leukocytosis, older school-age - macrolides should only be considered in true atypical pneumonia - If you are sure it is a type-1 reaction, can try cephalosporins (2nd or 3rd gen.) - Consider risk factors for MRSA CPS statement 2016

Community-acquired meningitis in children < 3 months (excluding neurosurgery or immunocompromised patients)	<i>Bacteroid</i> (<i>S. pneumoniae</i> , <i>N. meningitidis</i> , <i>H. influenzae</i>), <i>Viral</i> (HSV, Enterovirus) <i>Special considerations in:</i> - < 3mo - immunocompromised - known CNS disease, post-neurosurgery, trauma	Gefixaxone IV/IM (meningitic dose, 10mg/kg/day in 2 divided doses) PLUS Vancomycin 15mg/(kg·dose) IV q6h *above antibiotic choices may not apply to those with special considerations - ADD acyclovir if any of the following: significant change in LOC, seizures - MR or EEG consistent with HSV - HSV PCR positive	Depends on organism: <i>S. pneumoniae</i> 10-14 days <i>N. meningitidis</i> 5-7 days <i>H. influenzae</i> 7-10 days	Mandatory ID consult consider DEXAMETHASONE if bacterial pathogen suspected 0.6 mg/kg/day divided q6h before or within 30 minutes of the first dose of antibiotics (only continue for 2 days if <i>S. pneumoniae</i> or <i>H. influenzae</i> isolated, any other pathogen discontinue) <i>CPS statement 2014</i>
Urinary Tract Infection (> 2 months of age)	<i>E. coli</i> , <i>Klebsiella</i> , <i>Enterococcus</i> , <i>Proteus</i> , <i>Serratia</i> , <i>Pseudomonas</i> , <i>Staphylococcus saprophyticus</i> <i>Acronym: KEEPSS</i>	Albirelle UTI (cystitis) Cephalexin Trimethoprim/sulfamethoxazole IV/IM Tobramycin Uncomplicated febrile UTI (pyelonephritis): Trimethoprim/sulfamethoxazole (older children) Complicated (requires admission < 2 months, hemodynamically unstable, elevated serum creatinine, poor urinary flow, abdominal or bladder mass, vomiting, clinically deteriorating after 24 hours of appropriate antibiotics, immunocompromised): Ampicillin IV PLUS tobramycin IV	Albirelle UTI (cystitis): 5 days 5 days cephalexin trimethoprim/sulfamethoxazole x1 dose IV/IM (tobramycin) Febrile UTI (pyelonephritis): 7-10 days (usual duration) 7	- Diagnosis: urine analysis and culture (will only send culture if mid-stream, catheter or suprapubic aspiration i.e. NO BAG SAMPLES for culture) - UNLIKELY TO BE UTI IF URINALYSIS NORMAL in an immunocompetent patient (any age) - Urine analysis and urinalysis + urinalysis investigation with an abdominal ultrasound - AAP Practice Guideline 2011 - CPS Statement 2020
Celulitis	Group A <i>Streptococcus</i> , <i>S. aureus</i> (MSSA/MRSA), Group C/G <i>Streptococcus</i> If pus not present – very likely streptococcal	<u>Preferred:</u> 1 st gen cephs (Cephalexin PO/Cefazolin IV) <u>Unsuspect MRSA (eg. abscess, open DR, severe disease):</u> Trimethoprim/Sulfamethoxazole PO or Vancomycin IV if concerns of MRSA	5 days (usually 1-2 days after the rash resolves) Varies depending on presence of abscess and degree of drainage	- Must do I&D as first line if abscess or furuncle - Consider MRSA risk factors - avoid oral cloxacillin if possible as it has poor bioavailability and has GI side effects <i>AMM Practice Point 2022</i>
Bone and joint infection	Group A <i>Streptococcus</i> , <i>Staphylococcus aureus</i> , <i>Kingella kingae</i> (particularly in pre-school age), <i>Staphylococcus pneumoniae</i>	<u>Preferred:</u> 1 st gen cephalosporins (cefazolin IV) at 50mg/kg/DOSE IV q6h Vancomycin 20mg/kg/DOSE IV q6h or 15mg/kg/dose IV q6h and linzolid <u>Unsuspect MRSA:</u> Vancomycin 20mg/kg/DOSE IV q6h or 15mg/kg/dose IV q6h and linzolid Metronidazole 30mg/kg/DAY PO (or IV) TID or QID	In general, for acute uncomplicated infection, Septic arthritis 2-3 weeks Acute uncomplicated osteomyelitis 3-4 weeks General duration 6-10-14 days A course of vancomycin tapering may be considered in recurrent episodes	Mandatory ID consult <i>CPS Statement 2018</i>
<i>Clostridium difficile</i> infection (CDI) See algorithm below	<i>Clostridioides difficile</i> Mild to moderate Diarrhea BUT no systemic toxicity Severe disease Systemic toxicity +/- complications including hypotension, shock, toxic megacolon, severe colitis, ileus etc.	1 st episode (severe +/- complications) or recurrent disease Vancomycin 10mg/kg/DOSE QID (maximum 125mg/DOSE) *can consider rectal vancomycin if ileus present, see algorithm	- always reassess need for concomitant antibiotics - Don't send stool for <i>C. difficile</i> testing in children < 1 year of age - <i>C. difficile</i> testing should only be done on diarrheal stool - Do not send stool for test of cure - Strongly consider ID consult for severe CDI or recurrent disease	

Fever in a neonate (< 4 weeks) (presenting from home)	<p>Group B Streptococcus, gram negatives (<i>E. coli</i>), Enterococcus (<i>aureus</i>, <i>S. pneumoniae</i> less likely)</p> <p>HSV (usually before 4 weeks of age)</p> <p>Virus (e.g. Enterovirus)</p>	<p>If clinically stable and no concerns of meningitis: ampicillin + gentamicin</p> <p>If clinically unwell/septic: ampicillin + cefotaxime, consider acyclovir</p> <p>If meningitis suspected clinically (e.g. unwell, bulging fontanelle, seizures, posturing, significant lethargy) or CSF abnormalities, ensure that cefotaxime and acyclovir are given</p> <p>Empiric therapy: cefotaxime and reassess need for ongoing antibiotics in 24-36 hours</p>	<p>Duration will depend on final diagnosis</p>	<p>LP is usually warranted in a neonate who presented with a fever</p> <p>Indications for acyclovir not clear-cut. Should be given for any neonate with severe sepsis, especially if thrombocytopenia or transaminitis or coagulopathy is present, any neonate with CSF pleocytosis, or if vesicular rash. However, incidence of neonatal HSV disease low, most cases occur < 21 days.</p> <p>Any baby started on acyclovir requires at minimum:</p> <ol style="list-style-type: none"> 1. LP for HSV PCR 2. Mouth, rectal, conjunctival and vesicle swab for HSV PCR
Typhoid/Paratyphoid Fever	<p><i>Salmonella typhi</i> or <i>paratyphi</i></p> <p>Obtain blood culture in patients with fever with no clear focus and travel in past few weeks (max incubation period is 60day), or anyone with <i>Salmonella</i> positive stools</p> <p>Most commonly associated with GI symptoms, but can be CNS, MSK, disseminated.</p> <p>Assess for Malaria risk in anyone with fever and travel to a malaria endemic region - send malaria smear x 2-3 separated by 12-24 hours</p>	<p>If clinically unwell, <3months of age, asplenic - admit and start IV Ceftriaxone pending blood culture results.</p> <p>If clinically well, >3months - start Azithromycin, discuss with ID and ensure followup pending sensitivity results.</p>	<p>Uncomplicated typhoid/paratyphoid fever and PO stepdown:</p> <ul style="list-style-type: none"> • 7 days for azithromycin, • 3 to 7 days for quinolones, • 7 to 14 days for cefixime, • 10 to 14 days for ceftriaxone • 14 days for amoxicillin or TMP-SMX [9] 	<p>-stool cultures positive in 30% of patients</p> <p>-only use quinolones in lab confirmed sensitive strains due to rising resistance</p> <p>-relapse occurs in up to 17% of cases usually within 4 weeks</p> <p>-fever persists 6 to 8days from antibiotic start, and is not a contraindication to switch to PO antibiotics</p> <p>-counsel on pre travel vaccination for future</p>

CLINICAL PEARLS

Other Clinical Scenarios:	Challenging Organisms:	MRSA covered by:	Organisms resistant to penicillins and cephalosporins:	Antibiotics of note:	Carbapenem indications:
Septic Shock: <ul style="list-style-type: none"> - ceftriaxone + vancomycin - can consider pip-tazo if require coverage for anaerobes (eg. GI infection) or pseudomonas Febrile Neutropenia: <ul style="list-style-type: none"> - Piperacillin-tazobactam - Consider empiric vancomycin if previous infection/colonization with MRSA, or clinical signs of sepsis - Refine ABX if blood Cx - Consider previous microbiology history (eg. antibiotic-resistant organisms) - Please note that piperacillin-tazobactam does not have reliable CNS coverage 	Pseudomonas often covered by: <ul style="list-style-type: none"> - ceftazidime - piperacillin +/- tazobactam - ciprofloxacin - meropenem - aminoglycosides (gentamicin/tobramycin / amikacin) 	MRSA covered by: <ul style="list-style-type: none"> - Vancomycin - Septera - Clindamycin (increasing resistance) - Linezolid (needs ID endorsement) - Doxycycline (available as PO and generally not indicated unless > 8 years) Risk Factors: <ul style="list-style-type: none"> - Previous MRSA - Infection or household contact - Healthcare exposure/recent hospitalization - TRAVEL (including to USA) 	Organisms resistant to penicillins and cephalosporins: <ul style="list-style-type: none"> - MRSA - most CONS - ESKL - ECK (AmpC producers): <i>Enterobacter cloacae</i>, <i>Freundii</i>, <i>Klebsiella aerogenes</i> - Atypicals - C diff <p>Cephalosporins do not have activity against <i>Enterococcus</i> or <i>Listeria</i></p>	Antibiotics of note: <ul style="list-style-type: none"> - Vancomycin (only covers gram pos. infections) - MRSA - Severe c diff infection (PO only) - CONS - <i>Enterococcus</i> 	Carbapenem indications: <ul style="list-style-type: none"> - ESBL - ECK (AmpC producers): <i>Enterobacter cloacae</i>, <i>Citrobacter freundii</i>, <i>Klebsiella aerogenes</i> - Polymicrobial CNS infection <p>REQUIRES ID CONSULT</p>



Pediatric Blood Culture Guidelines

1. **AEROBIC** cultures are always drawn.

2. Does the patient require an **ANAEROBIC** culture as well?

☐ **YES** if greater than 45kg

☐ **YES** if less than 45 kg **AND** if any of these conditions are suspected



Specimen Labels: Position lengthwise ensuring QR code and specimen window are not covered

1. Intra-abdominal or pelvic infection
2. NEC or Intestinal perforation in a neonate
3. Necrotizing soft tissue infection (e.g. Necrotizing fasciitis)
4. Chronic oral or sinus infections with sepsis
5. Infected Bite Wound
6. Deep neck space infections (e.g. Lemierre's Syndrome)
7. Immunocompromised (e.g. Febrile neutropenic)
8. Prolonged fever of unknown origin with negative aerobic culture

For Peripheral cultures only: If patient has a Central Vascular Access Device, see instructions and chart on reverse.

Peripheral blood culture requirements:

- Find patient weight on chart below to see total volume of blood required
- Look at the appropriate section, **aerobic only** or **aerobic + anaerobic** to see how the total volume is divided. (Number of bottles, bottle colour and volume)
- Blood is collected from one peripheral poke, unless > 45kg
- If unable to obtain required blood volume, refer to min and max blood volume reference and adjust as needed
- For patients >45kg, if unable to get a 2nd site after the most proficient RN attempt, notify MRP for further direction

Min and Max Blood Volumes per Bottle Type		
Aerobic Bottles		Anaerobic Bottle
Yellow 1.5 – 5.0mL	Green 5.1-10mL	Orange 1 - 10 mL

Weight	Total Blood Volume	Aerobic		Aerobic + Anaerobic	
		Volume per bottle and required bottle(s)	Volume per bottle and required bottle(s)	Volume per bottle and required bottle(s)	Volume per bottle and required bottle(s)
< 5 kg	2-4 mL divided →	2 – 4mL in yellow	1.5 – 2mL in yellow	1.5 – 2mL in yellow	1.5 – 2mL in orange
5 – 13 kg	5-7 mL divided →	5 – 7mL (yellow if 5, otherwise green)	2.5 – 3.5mL in yellow	2.5 – 3.5mL in yellow	2.5 – 3.5mL in orange
13 – 36 kg	14-20 mL divided →	7 – 10mL in green 7 – 10mL in green	10 mL in green 5 – 10mL (yellow if 5, otherwise green)	7 – 10mL in green	7 – 10mL in orange
36 – 45 kg	21-30 mL divided →	10mL in green 5 – 10mL (yellow if 5, otherwise green)	6 – 10mL in green	10 mL in green 5 – 10mL (yellow if 5, otherwise green)	6 – 10mL in orange
> 45 kg	40 mL divided →	Aerobic and Anaerobic always drawn. Requires sample from 2 separate sites.		Site 1: 10mL in green 10mL in orange	Site 2: 10mL in green 10mL in orange

Minimize contamination during collection:

- Disinfect skin with a chlorhexidine with a contact time of 30 seconds
- Disinfect septum of BC bottle with alcohol pad for 15 seconds
- Do not re-palpate skin

For Central and Peripheral cultures: If Patient has a Central Vascular Access Device (CVAD)

Central line (CVAD) blood culture requirements

- Cultures from CVAD and peripheral draw required. A peripheral culture is essential to guide management.
- If the most proficient provider has tried and is unable to obtain a peripheral culture, a second CVAD culture can be done with a new set-up
- If CVAD has multiple lumens, all lumens must be cultured (sometimes the bug will only be found in 1 of the lumens)
- If anaerobic culture indicated, and patient is < 45 kg, only need 1 anaerobic sample from any site

Use chart below to determine number of sample sites, blood volumes and bottles required

- Identify number of lumens on CVAD, type of culture (aerobic or aerobic + anaerobic) and weight of patient on chart
- Draw volume of blood listed from each site indicated and place in coloured bottle(s) noted

Legend:		P= Peripheral	L1= CVAD Lumen #1	L2= Double lumen CVAD	L3= CVAD Lumen #2	L3= CVAD Lumen #3
		Single lumen CVAD		Double lumen CVAD		Triple lumen CVAD
Weight	Total blood volume	Aerobic	Aerobic + Anaerobic	Aerobic	Aerobic + Anaerobic	Aerobic Aerobic + Anaerobic
< 5 kg	2-4 mL divided →	L1: 1.5-2 in yellow P: 1.5-2 in yellow	L1: 1.5 in yellow 1.5 in orange P: 1.5 in yellow	L1: 1.5 in yellow L2: 1.5 in yellow P: 1.5 in yellow	L1: 1.5 in yellow 1 in orange L2: 1.5 in yellow P: 1.5 in yellow	Refer to Neo policy
5 – 13 kg	5-7 mL divided →	L1: 2.5-3.5 in yellow P: 2.5-3.5 in yellow	L1: 1.5-3 in yellow 2 in orange P: 1.5-2 in yellow	L1: 1.5-3 in yellow L2: 2 in yellow P: 1.5-2 in yellow	L1: 1.5-2 yellow 1-2 orange L2: 1.5 yellow P: 1.5 yellow	L1: 1.5 yellow 1 orange L2: 1.5 yellow L3: 1.5 yellow P: 1.5 yellow
13.1 – 36 kg	14-20 mL divided →	L1: 7-10 green P: 7-10 green	L1: 5-10 (yellow if 5, otherwise green) 5 in orange P: 4-5 in yellow	L1: 5-10 (yellow if 5, otherwise green) L2: 5 in yellow P: 4-5 in yellow	L1: 4-5 yellow L2: 4-5 yellow L3: 4-5 yellow P: 2-5 yellow	L1: 5 yellow 5 orange L2: 1.5-4 yellow L3: 1.5-4 yellow P: 1.5-2 yellow
36.1 – 45 kg	21-30 mL divided →	L1: 10 green 6-10 green P: 5-10 (yellow if 5, otherwise green)	L1: 10 green 6-10 orange P: 5-10 (yellow if 5, otherwise green)	L1: 10 green L2: 6-10 green P: 5-10 (yellow if 5, otherwise green)	L1: 6-10 green 5-10 orange L2: 5 yellow P: 5 yellow	L1: 5-10 (yellow if 5, otherwise green) 5 orange L2: 5 yellow L3: 4-5 yellow P: 2-5 yellow
> 45 kg	40 mL divided →	Aerobic and Anaerobic always drawn.	L1: 10 green 10 orange P: 10 green 10 orange	Aerobic and Anaerobic always drawn.	L1: 10 green 10 orange P: 10 green 5 orange	L1: 10 green 10 orange L2: 5 yellow L3: 5 yellow P: 5 yellow 5 orange

CONSIDERATION OF MULTISYSTEM INFLAMMATORY SYNDROMES, INCLUDING MULTISYSTEM INFLAMMATORY SYNDROME IN CHILDREN ASSOCIATED WITH COVID-19 (MIS-C) AND KAWASAKI DISEASE (KD)			Toxic Shock Syndrome
Clinical features	Fever >38.5 °Celsius of any duration AND Tachycardia, hypotension and/or oxygen requirement AND Evidence of multi-organ dysfunction	Persistent fever >38.5 °Celsius for 5 days * plus/minus Additional Clinical Features of KD and MIS-C (See Boxes A and B)	Staph: Fever, hypotension, diffuse erythroderma, desquamation, at least 3 organ systems No other infection confirmed Group A Strep: -hypotension, multiorgan involvement (renal, coagulopathy, liver, ARDS, erythematous macular rash with desquamation) with no other etiology. -isolation of GAS from sterile site
Initial Action Steps	Consider early PICU consult Order MIS-C Full Panel (See Box C)	Order MIS-C Screening Blood Work (See Box D)	-send blood cultures (two sets) -culture sterile sites of relevance (e.g. joints, wounds, CNS etc as appropriate) -remove any source foreign bodies (e.g. tampons)
Assessment of MIS-C Screen		Are at least 2 of the following present?	
Next Action Steps		1. CBC abnormalities consistent with MIS-C (e.g. neutrophilia, lymphopenia, anemia or thrombocytopenia) 2. CRP > 100 3. Ferritin > 500 4. Albumin < 30	
		YES	NO
	Contact General Pediatrics and Rheumatology to discuss potential admission and/or further work-up	Manage as per usual care by the ED team	-initiate antibiotics Ceftriaxone empirically (if MRSA risk factors consider Vancomycin) -in addition to Clindamycin (30-40mgkg/day IV divided q8h) Consider IVIG in GAS TSS (1g/kg on day 1, 0.5g/kg on day 2, 3) Consider prophylaxis for close contacts of iGAS
* Physicians should use their clinical judgement:			
<ul style="list-style-type: none">Patients with an obvious cause for their fever (e.g. Streptococcal pharyngitis or pneumonia) <u>may not require</u> MIS-C screening blood workPatients with less than 5 days of fever and 2 or more concerning symptoms for MIS-C (see Box A) may require screening blood work			

Box A. Additional clinical features of Multisystem Inflammatory Syndrome in Children (MIS-C)

- Abdominal pain, diarrhea and/or vomiting
- Non-exudative conjunctivitis
- Oral mucosal changes
- Hand and foot erythema or edema
- Diffuse erythematous rash
- Headache or neck stiffness

Box B. Features that help distinguish patients with MIS-C from those with Kawasaki disease (KD)

	KD	MIS-C
Age	• Typically 1 to 5 years of age	• Often older than typical KD patients
Clinical features	<ul style="list-style-type: none"> • Classic KD features present <ul style="list-style-type: none"> ◦ Nonpurulent conjunctivitis ◦ Oral mucosal changes ◦ Cervical lymphadenopathy ◦ Rash ◦ Peripheral edema 	<ul style="list-style-type: none"> • Classic KD features present, but more likely to be incomplete • More respiratory and gastrointestinal symptoms • Meningeal signs may be present • More likely to have signs of cardiovascular involvement
Laboratory features	<ul style="list-style-type: none"> • Leukophilia • Normal/increased lymphocytes • Increased platelet counts • Mild to moderately high ferritin • Normal CK 	<ul style="list-style-type: none"> • May have leukopenia • Lymphopenia • Thrombocytopenia • Higher ferritin levels • Elevated CK
Clinical course		<ul style="list-style-type: none"> • More severe disease course • More likely to have myocarditis and shock • More IVIG resistance • Increased rates of cytokine storm (e.g. secondary HLH)

Box C. MIS-C Full Panel

- CBC including differential, blood film
- CRP
- Ferritin
- Albumin
- ALT, AST, GGT, LDH, bilirubin
- Electrolytes, urea, creatinine
- Glucose
- Blood gas
- INR, PTT, fibrinogen, D-Dimer
- Triglycerides
- CK, troponin, BNP (or NT-pro-BNP)
- Save full red top tube for COVID-19 serological testing prior to IVIG
- Blood culture
- NPS for respiratory viruses plus SARS-CoV-2
- Urinalysis

Box D. MIS-C Screening Blood Work

- CBC including differential
- CRP
- Ferritin
- Albumin
- ALT
- Creatinine

APPENDIX A

Centers for Disease Control and Prevention Case Definition of Multisystem Inflammatory Syndrome in Children (MIS-C) Associated with Coronavirus Disease 2019 (COVID-19)

- An individual aged <21 years presenting with feverⁱ, laboratory evidence of inflammationⁱⁱ, and evidence of clinically severe illness requiring hospitalization, with multisystem (≥2) organ involvement (cardiac, renal, respiratory, hematologic, gastrointestinal, dermatologic or neurological); **AND**
- No alternative plausible diagnoses; **AND**
- Positive for current or recent SARS-CoV-2 infection by RT-PCR, serology, or antigen test; or COVID-19 exposure within the 4 weeks prior to the onset of symptoms

ⁱ Fever ≥38.0°C for ≥24 hours, or report of subjective fever lasting ≥24 hours

ⁱⁱ Including, but not limited to, one or more of the following: an elevated C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), fibrinogen, procalcitonin, d-dimer, ferritin, lactic acid dehydrogenase (LDH), or interleukin 6 (IL-6), elevated neutrophils, reduced lymphocytes and low albumin

Additional comments

- *Some individuals may fulfill full or partial criteria for Kawasaki disease but should be reported if they meet the case definition for MIS-C*
- *Consider MIS-C in any pediatric death with evidence of SARS-CoV-2 infection*

Source: <https://emergency.cdc.gov/han/2020/han00432.asp>

APPENDIX B

World Health Organization Preliminary Case Definition of Multisystem Inflammatory Syndrome in Children and Adolescents Temporally Associated with COVID-19

Children and adolescents 0–19 years of age with fever ≥ 3 days

AND 2 of the following:

1. Rash or bilateral non-purulent conjunctivitis or muco-cutaneous inflammation signs (oral, hands or feet).
2. Hypotension or shock.
3. Features of myocardial dysfunction, pericarditis, valvulitis, or coronary abnormalities (including ECHO findings or elevated Troponin/NT-proBNP),
4. Evidence of coagulopathy (by PT, PTT, elevated d-Dimers).
5. Acute gastrointestinal problems (diarrhoea, vomiting, or abdominal pain).

AND

Elevated markers of inflammation such as ESR, C-reactive protein, or procalcitonin.

AND

No other obvious microbial cause of inflammation, including bacterial sepsis, staphylococcal or streptococcal shock syndromes.

AND

Evidence of COVID-19 (RT-PCR, antigen test or serology positive), or likely contact with patients with COVID-19.

Source: <https://www.who.int/news-room/commentaries/detail/multisystem-inflammatory-syndrome-in-children-and-adolescents-with-covid-19>

APPENDIX C

Proposed updated definition - definitive case of MIS-C/A (Level 1 of Diagnostic Certainty)

Age < 21 years (MIS-C) OR > 21 years (MIS-A)

AND fever for ≥ 3 consecutive days

AND 2 or more of the following clinical features:

- Mucocutaneous (rash, erythema or cracking of the lips/mouth/pharynx, bilateral nonexudative conjunctivitis, erythema/edema of hands and feet)
- Gastrointestinal (abdominal pain, vomiting, diarrhea)
- Shock or hypotension
- Neurologic (altered mental status, headache, weakness, paresthesias, lethargy)

AND laboratory evidence of inflammation including any of the following:

- Elevated CRP
- Elevated ESR
- Elevated ferritin
- Elevated procalcitonin

AND 2 or more measures of disease activity:

- Elevated BNP, NT-proBNP or troponin
- Neutrophilia, lymphopenia, or thrombocytopenia
- Evidence of cardiac involvement by echocardiography (dysfunction, wall motion abnormality, coronary abnormality, valvular regurgitation, pericardial effusion, abnormal LV strain) or physical stigmata of heart failure (gallop, rales, lower extremity edema, hepatosplenomegaly, jugular venous distension)
- ECG changes consistent with myocarditis or myopericarditis (abnormal ST segments, arrhythmia, pathogenic Q waves, AV conduction delay, PR segment depression, low voltage QRS)

AND

- Laboratory confirmed SARS-CoV-2 infection
- Personal history of confirmed COVID-19 within 12 weeks
- Close contact with known COVID-19 case within 12 weeks
- Following SARS-CoV-2 vaccination

Source: Vogel TP et al. Multisystem inflammatory syndrome in children and adults (MIS-C/A): Case definition & guidelines for data collection, analysis, and presentation of immunization safety data. *Vaccine* 2021; 39:3037-49.

Reportable diseases to public health

- [Acquired Immunodeficiency Syndrome \(AIDS\)](#)
- Acute Flaccid Paralysis
- Amebiasis
- Anthrax
- Blastomycosis
- Botulism
- Brucellosis
- Campylobacter enteritis
- Carbapenemase-producing Enterobacteriaceae (CPE), infection or colonization
- Chancroid
- [Chickenpox \(Varicella\)](#)
- [Chlamydia trachomatis infections](#)
- Cholera
- Clostridium difficile infection (CDI) outbreaks in public hospitals
- Creutzfeldt-Jakob Disease
- Cryptosporidiosis
- Cyclosporiasis
- Diphtheria
- Diseases caused by a novel coronavirus, including Severe Acute Respiratory Syndrome (SARS), Middle East Respiratory Syndrome (MERS) and coronavirus disease (COVID-19).
- Echinococcus multilocularis infection
- Encephalitis, including,
 - i. [Primary, viral](#)
 - ii. Post-infectious
 - iii. Vaccine-related
 - iv. Subacute sclerosing panencephalitis
 - v. Unspecified
- Food poisoning, all causes
- Gastroenteritis, outbreaks in institutions and public hospitals
- Giardiasis, except asymptomatic cases
- [Gonorrhoea](#)
- [Group A Streptococcal disease, invasive](#)
- Group B Streptococcal disease, neonatal
- Haemophilus influenza disease, all types, invasive
- Hantavirus pulmonary syndrome
- Hemorrhagic fevers, including,
 - i. Ebola virus disease
 - ii. Marburg virus disease
 - iii. Lassa fever, and other viral causes
- [Hepatitis A, viral](#)
- [Hepatitis B, viral](#)
- [Hepatitis C, viral](#)
- Influenza
- [Legionellosis](#)
- Leprosy
- Listeriosis
- [Lyme Disease](#)
- [Measles](#)
- Meningitis, acute,
 - i. [bacterial](#)
 - ii. [viral](#)
 - iii. [other](#)
- [Meningococcal disease, invasive](#)
- [Mumps](#)
- Ophthalmia neonatorum
- Paralytic Shellfish Poisoning
- Paratyphoid Fever
- [Pertussis \(Whooping Cough\)](#)
- Plague
- Pneumococcal disease, invasive
- Poliomyelitis, acute
- Psittacosis/Ornithosis
- Q Fever
- [Rabies](#)
- Respiratory infection outbreaks in institutions and public hospitals
- Rubella
- Rubella, congenital syndrome
- Salmonellosis
- Shigellosis
- Smallpox and other orthopoxviruses, including monkeypox
- [Syphilis](#)
- Tetanus
- Trichinosis
- [Tuberculosis](#)
- Tularemia
- Typhoid Fever
- Verotoxin-producing E. coli infection, including Haemolytic Uremic Syndrome (HUS)
- [West Nile Virus Illness](#)
- Yersiniosis

For specifics on which diseases need to be reported immediately, same day and next business day visit: <https://www.hamilton.ca/people-programs/public-health/health-care-professionals/reporting-infectious-diseases#report-same-day>

How to report:

Timely reporting can help minimize the spread of communicable diseases. If you suspect, or have confirmation of, any of the below named diseases or their etiologic agent: Contact Hamilton Public Health by phone at [905-546-2063](tel:905-546-2063) or by fax at 905-546-4078

Post Exposure Prophylaxis for specific pathogens:

Pathogen	Who to Prophylax	Agents of Prophylaxis	Timing
N. meningitidis	<ul style="list-style-type: none"> -all household contacts -shared utensils -slept in same dwelling 7d prior -anyone with contact with oral/nasal secretions -inform public health if index was on a flight in the 7d prior -specific high risk healthcare workers (EHS responsibility) 	<p>Rifampin PO x 2 days (preferred agent for all infants) (consider drug interactions for adults)- not to use in pregnancy</p> <p>Ceftriaxone IM x 1</p> <p>Ciprofloxacin x 1 dose</p> <p>-Vaccinate any household contacts that are not vaccinated and eligible</p>	<p>Initiate within 24h after index case identified and within 2 weeks</p> <p>https://www.health.gov.on.ca/en/pro/programs/publichealth/oph_standards/docs/meningococcal_chapter.pdf</p>
H. influenzae B	<ul style="list-style-type: none"> -household contacts if any children <4y of age in home (if not fully vaccinated) or any unvaccinated children or immune compromised child in the home -daycare contacts if incompletely immunized contacts 	<p>Rifampin x 4 days</p> <p>-Vaccinate any household contacts/exposed people that are not vaccinated</p>	<p>Initiate within 7days of index case</p> <p>https://www.health.gov.on.ca/en/pro/programs/publichealth/oph_standards/docs/Haemophilus_influenzae_chapter.pdf</p>
Invasive Group A Strep	<ul style="list-style-type: none"> -household contacts (e.g. >4h / days or >20h/ week in the past 7days) -shared same bed -anyone with contact with oral/nasal secretions or shared needles -specific high risk healthcare workers (EHS responsibility) 	<p>Cephalexin X 10days</p> <p>Clarithromycin x 10days</p> <p>Clindamycin x 10days</p>	<p>Initiate within 7days of index case</p> <p>https://cps.ca/en/documents/position/invasive-group-A-streptococcal-disease</p>

Drug Generic Name	Brand Name	Pediatric Dose ⁶ (<i>BID dosing is thought to provide better control of acid</i>)	Max Dose ⁶ (faster clearance in peds than adults – may need higher than standard adult dose)	Usual Adult Dose GERD ²	Administration (See note below) Note: <i>Pharmacy Prepared Suspension⁸ (Compounding dependent on pharmacy)</i>	Available Formats ³ and Cost	LU Code ³
Omeprazole	Losec	1-1.5 mg/kg/day PO once daily or divided BID NEONATAL: 0.5-1.5 mg/kg/dose	3.5 mg/kg/day	10-20 mg PO OD	1. Capsule – can be opened & sprinkled on yogurt and given 2. Pharmacy prepared suspension can be used	10mg capsules – not ODB covered 20 mg cap (\$0.41/cap)	293 – GERD or non erosive GERD when H ₂ Antags have failed 297-PUD or prevention of NSAID induced ulcers 401- treatment of GI disorders, Crohns, short Gut etc. 402-severe esophagitis, Zollinger-Ellison etc.
Lansoprazole	Prevacid	<10 kg: 7.5 mg PO OD 10-30 kg: 15 mg PO OD >30 kg: 30 mg PO OD	1.6 mg/kg/day or 30 mg/day	15-30 mg PO OD	1. Capsules may be opened and sprinkled into applesauce 2. Fast tabs can be placed on tongue for doses 15mg or greater 3. Fast tabs can be split and mixed with water if no other options exist (cannot dissolve and dose) 4. Pharmacy Prepared suspension has short expiry so not made at HHS	15mg (\$0.5/cap) 30mg (\$0.5/cap) with Enteric coated microgranules 15, 30 mg Fast tabs (not ODB covered)	Required for billing of suspension For capsules only: (not Fast tabs) 293 – GERD or non erosive GERD when H ₂ Antags have failed 295 – for HPYlon Pepsic Ulcer 297-PUD or prevention of NSAID induced ulcers 401- treatment of GI disorders, Crohns, short Gut etc. 402-severe esophagitis, Zollinger-Ellison etc.
Esomeprazole	Nexium	1mg-11 yrs: -5kg: 2.5-5mg PO OD 5-10kg: 5-10mg PO OD 10-17yrs: 20 mg PO OD	40 mg/day	20-40 mg PO OD	1. Tabs can be dispersed for PO admin. Mix with 25-50mL mL of water. 2. Sachet can be dissolved & administered via G tube	20 mg, 40 mg tablets (\$0.36/40mg tab) 10 mg sachet for oral suspension (Not ODB covered)	NO – Not covered under ODB
Pantoprazole	Pantoloc	1-1.5 mg/kg/day	40 mg/dose	20-40 mg PO OD	Cannot be crushed	20mg- not a benefit 40 mg (\$0.3/tablet)	See above (same as omeprazole)
Rabeprazole	Pariet	Greater than 10 years: 10 mg PO OD		20 mg PO OD	Cannot be crushed	10 mg (\$0.12 tablet), 20 mg (\$0.24/tablet)	No LU code required

Note: Directions for opening capsules and dissolving tablets with dispersed microgranules into food or water requires that the granules must NOT be crushed or chewed for effect.

- Hospital for Sick Children, Drug Handbook and Formulary, 2016.
- RX Files Drug Comparison Charts, 8th Edition
- ODB Drug Formulary
- eCPS, 2016
- Jaw, RK et al. Extemporaneous Formulations for Pediatric, Geriatric, and Special Needs Patients. ASHP, 2nd Edition.
- Micromedex. Accessed May 2017.

Prepared by N Fernandes RPh, Drug Information Centre, HHS. Reviewed by N Clarke RPh, Pediatrics MCH.

Escalation Protocol

Continue plan of care
<ul style="list-style-type: none"> Initiate nursing directed interventions* Within 90 minutes post intervention(s), repeat HPEWS vital signs. If patient remains YELLOW, notify charge RN, medical team and RT if applicable
In addition to nursing directed interventions: <ul style="list-style-type: none"> NOTIFY charge RN, medical team (team resident**/fellow/NP) and RT if applicable Within 60 minutes post interventions, repeat HPEWS vital signs. If patient remains ORANGE, re-notify team as above
In addition to nursing directed interventions: <ul style="list-style-type: none"> NOTIFY charge RN, medical team (team resident**/fellow/NP) and activate PACE Within 30 minutes post interventions, repeat HPEWS vital signs If patient remains RED, re-notify team as above

*nursing directed interventions include increased frequency of vital signs, repositioning, comfort measures, prn medications etc.

**junior residents must review with senior resident

At any time regardless of HPEWS colour, anyone can notify/activate MRP team, PACE team, RT or Pediatric Code Blue



Pediatric Assessment of Critical Events (PACE) PACE Calling Criteria

Call PACE in the following situations:

If the health care provider or family member is worried about the patient's clinical state or if any of the following criteria are present

Airway

Threatened or obstructive symptoms: stridor, excessive secretions

Breathing

Severe respiratory distress, apnea, tachypnea or cyanosis

Age	Respiratory rate/min	Hypoxemia
Term – 3 months	> 60	SaO ₂ < 90% in >40% FiO ₂ SaO ₂ < 60% in > 40% FiO ₂ (cyanotic heart disease)
4-12 months	> 50	
1-4 years	> 40	
5-12 years	> 30	
12 years +	> 30	

Circulation

Age	Bradycardia (beats/min)	Tachycardia (beats/min)	BP (systolic mmHg)
Term – 3 months	< 100	> 180	< 50
4-12 months	< 100	> 180	< 60
1-4 years	< 90	> 160	< 70
5-12 years	< 80	> 140	< 80
12 years +	< 60	> 130	< 90

Neurologic State

Acute change in neurologic status or convulsion


- Some of the values for respiratory rate, heart rate and blood pressure are outside the normal ranges for age: they represent concerning levels that may indicate serious illness and require expert review
- It is also important to look for worsening trends in vital signs and report these.

**Call ext. 75030 and ask for PACE.
We're here to help!**

During the training phase, the PACE team will be available from Monday to Friday, 8 a.m. to 4 p.m.
Coming soon! On January 29th, 2007 we will begin providing 24-hour daily coverage.

Activate 'Code Blue' for all respiratory and/or cardiac arrests
or other medical emergencies as per HHS policy.

Pediatric Cardiac Arrest Algorithm



CPR Quality

- Push hard ($\geq\frac{1}{2}$ of anteroposterior diameter of chest) and fast (100-120/min) and allow complete chest recoil
- Minimize interruptions in compressions
- Change compressor every 2 minutes, or sooner if fatigued
- If no advanced airway, 15:2 compression-ventilation ratio
- If advanced airway, provide continuous compressions and give a breath every 2-3 seconds

Shock Energy for Defibrillation

- First shock 2 J/kg
- Second shock 4 J/kg
- Subsequent shocks ≥ 4 J/kg, maximum 10 J/kg or adult dose

Drug Therapy

- Epinephrine IV/IO dose:** 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration). Max dose 1 mg. Repeat every 3-5 minutes. If no IV/IO access, may give endotracheal dose: 0.1 mg/kg (0.1 mL/kg of the 1 mg/mL concentration).
- Amiodarone IV/IO dose:** 5 mg/kg bolus during cardiac arrest. May repeat up to 3 total doses for refractory VF/pulseless VT or
- Lidocaine IV/IO dose:** Initial: 1 mg/kg loading dose


Advanced Airway

- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement


Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypoglycemia
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary

Pediatric Basic Life Support Algorithm for Healthcare Providers—Single Rescuer



Pediatric Bradycardia With a Pulse Algorithm



© 2020 American Heart Association

Pediatric Tachycardia With a Pulse Algorithm

